

Clinical Policy: Celecoxib (Celebrex, Elyxyb)

Reference Number: IL.PMN.122

Effective Date: 6.26.23

Last Review Date: 6.26.23

Line of Business: Youthcare Healthchoice Illinois

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Celecoxib (Celebrex[®], Elyxyb[™]) is a nonsteroidal anti-inflammatory drug (NSAID).

FDA Approved Indication(s)

Celebrex is indicated for the treatment of:

- Osteoarthritis
- Rheumatoid arthritis
- Juvenile rheumatoid arthritis in patients 2 years and older
- Ankylosing spondylitis
- Acute pain
- Primary dysmenorrhea

Elyxyb is indicated for the acute treatment of migraine with or without aura in adults.

Limitation of use: Elyxyb is not indicated for the preventive treatment of migraine.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Celebrex and Elyxyb are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Acute Migraine Treatment (must meet all):

1. Request is for Elyxb;
2. Diagnosis of migraine;
3. Age \geq 18 years;
4. Member meets one of the following (a, b, c, d, or e):
 - a. Age $>$ 65 years;
 - b. Current use of a corticosteroid;
 - c. Current use of an anticoagulant or antiplatelet (e.g., aspirin, warfarin, low molecular weight heparin, direct thrombin inhibitors, factor Xa inhibitors, clopidogrel);
 - d. Prior gastrointestinal bleed or active peptic ulcer disease (not gastroesophageal reflux disease [GERD]);

- e. Failure of a ≥ 4 week trial of both of the following (i and ii), up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced:
 - i. Meloxicam;
 - ii. One additional generic NSAID;
5. Dose does not exceed 120 mg (4.8 mL) per day.

Approval duration: – 12 months

B. All Other Indications – (must meet all):

1. Request is for Celebrex;
2. Age ≥ 2 years;
3. For brand Celebrex requests, member must use generic celecoxib, unless contraindicated or clinically significant adverse effects are experienced;
4. Dose does not exceed both of the following (a and b):
 - a. 800 mg per day;
 - b. 2 capsules per day.

Approval duration: 12 months

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Acute Migraine Treatment (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed 120 mg (4.8 mL) per day.

B. All Other Indications – (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed both of the following (i and ii):
 - a. 800 mg per day;
 - b. 2 capsules per day.

Approval duration: 12 months

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CABG: coronary artery bypass graft

FDA: Food and Drug Administration

GERD: gastroesophageal reflux disease

NSAID: nonsteroidal anti-inflammatory drug

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
naproxen sodium (Anaprox [®] , Anaprox DS [®])	275 – 550 mg PO BID	1,650 mg/day
sulindac (Clinoril [®])	150 – 200 mg PO BID	400 mg/day
salsalate (Disalcid [®])	500 – 750 mg PO TID, titrated up to 3,000 mg/day	3,000 mg/day
piroxicam (Feldene [®])	10 – 20 mg PO QD	20 mg/day
indomethacin (Indocin [®])	25 – 50 mg PO BID -TID	200 mg/day
indomethacin SR (Indocin [®] SR)	75 mg PO QD - BID	150 mg/day
meclofenamate (Meclomen [®])	50 – 100 mg PO Q4-6hr	400 mg/day
meloxicam (Mobic [®])	7.5 – 15 mg PO QD	15 mg/day
ibuprofen (Motrin [®])	400 – 800 mg PO Q6-8hr	3,200 mg/day
fenoprofen (Nalfon [®])	200 mg PO Q4-6hr	3,200 mg/day
naproxen (Naprosyn [®])	250 – 500 mg PO BID	1,500 mg/day
ketoprofen (Orudis [®])	25 – 75 mg PO Q6-8hr	300 mg/day
nabumetone (Relafen [®])	1000 mg PO QD or 500 mg PO BID	2,000 mg/day
tolmetin (Tolmetin [®] DS)	400 mg PO TID, titrated up to 1800 mg/day	1,800 mg/day
diclofenac sodium (Voltaren [®])	50 mg PO Q6-8hr	200 mg/day
oxaprozin (Daypro [®])	600 – 1,200 mg PO QD	1,800 mg/day
etodolac (Lodine [®])	400 – 500 mg PO BID	1,200 mg/day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity to celecoxib or any components of the drug product; history of asthma, urticaria, or other allergic-type reactions to aspirin or other NSAIDs; in the setting of coronary artery bypass graft (CABG) surgery; allergic-type reactions to sulfonamides
- Boxed warning(s): increased risk of serious cardiovascular thrombotic events, including myocardial infarction, and stroke; increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines; in the setting of CABG surgery

Appendix D: General Information

- The risk vs. benefit of COX-II therapy should be individualized based on patient's previous GI history, other co-morbid conditions (e.g., angina, ischemic heart disease,

myocardial infarction (MI), coronary artery disease, stroke), age, concurrent medications (e.g., warfarin, oral corticosteroids), duration and dose.

- Celebrex has been associated with an increased risk of serious adverse cardiovascular (CV) events in a long-term placebo-controlled trial. Based on the currently available data, FDA has concluded that an increased risk of serious adverse CV events appears to be a class effect of NSAIDs. FDA has requested that the package insert for all NSAIDs, including Celebrex, be revised to include a boxed warning to highlight the potential increased risk of CV events and the well described risk of serious, and potentially life-threatening, gastrointestinal bleeding.

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Celecoxib (Celebrex)	Osteoarthritis	200 mg PO QD or 100 mg PO BID	800 mg/day
	Rheumatoid arthritis	100 to 200 mg PO BID	800 mg/day
	Juvenile rheumatoid arthritis	10-25 kg: 50 mg PO BID > 25 kg: 100 mg PO BID	200 mg/day
	Ankylosing spondylitis	200 mg PO QD or 100 mg PO BID. If no effect is observed after 6 weeks, a trial of 400 mg (single or divided doses) may be of benefit.	800 mg/day
	Acute pain or Primary dysmenorrhea	400 mg PO initially, followed by a 200 mg dose if needed on the first day. On subsequent days, 200 mg PO BID as needed	800 mg/day
Celecoxib (Elyxyb)	Migraine	120 mg PO PRN. Use the fewest number of days per month, as needed.	120 mg/day

VI. Product Availability

Drug Name	Availability
Celecoxib (Celebrex)	Capsules: 50 mg, 100 mg, 200 mg, and 400 mg
Celecoxib (Elyxyb)	Oral solution: 120 mg/4.8 mL (25 mg/mL)

VII. References

1. Celebrex Prescribing Information. New York, NY: G.D. Searle, LLC; May 2019. Available at: <http://www.celebrex.com/>. Accessed January 31, 2023.
2. Elyxyb Prescribing Information. Hyderabad, Telangana India: Dr. Reddy's Laboratories Limited; August 2021. Available at: www.elyxyb.com. Accessed January 31, 2023.
3. Lanza FL, Chan FK, Quigley EM et al. Guidelines for prevention of NSAID-related ulcer complications. Am J Gastroenterol. 2009 Mar;104(3):728-38. doi: 10.1038/ajg.2009.115. Epub 2009 Feb 24.

4. Kolasinski SL, Neogi T, Hochberg MC et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the management of osteoarthritis of the hand, hip, and knee. *Arthritis & Rheumatology*. 2020 Feb; 72(2): 220-233.
5. Fraenkel L, Bathon JM, England BR et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Care & Research*. 2021 July; 73(7):924-939.
6. Onel KB, Horton DB, Lovell DJ et al. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. *Arthritis & Rheumatology*. 2022 April; 74(4):553-569.
7. Ringold S, Weiss PF, Beukelman T et al. 2013 update of the 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: recommendations for the medical therapy of children with systemic juvenile idiopathic arthritis and tuberculosis screening among children receiving biologic medications. *Arthritis Rheum*. 2013 Oct;65(10):2499-512. doi: 10.1002/art.38092.
8. Ringold S, Angeles-Han ST, Beukelman T et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the treatment of idiopathic arthritis: therapeutic approaches for non-systemic polyarthritis, sacroiliitis, and enthesitis. *Arthritis Care & Research*. 2019 June; 71(6): 717-734. doi: 10.1002/acr.23870.
9. Ware MM, Deodhar A, Akl EA et al. American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network 2015 Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol*. 2016 Feb;68(2):282-98. doi: 10.1002/art.39298.
10. Yeomans ND. A comparison of omeprazole with ranitidine for ulcers associated with nonsteroidal anti-inflammatory drugs. *N Engl J Med* 1998;338:727-734.
11. Silverstein, et al. Gastrointestinal toxicity with celecoxib vs. nonsteroidal antiinflammatory drugs for osteoarthritis and rheumatoid arthritis (CLASS Study). *JAMA* 2000;284:1247-1255.
12. Mukherjee, et al. Risk of cardiovascular events associated with selective COX-2 inhibitors. *JAMA* 2001;286:954-959.
13. Juni, et al. Are selective COX 2 inhibitors superior to traditional non-steroidal anti-inflammatory drugs. *BMJ* 2002;324:1287-1288.
14. Solomon DH, et al. Relationship between selective cyclooxygenase-2 inhibitors and acute myocardial infarction in older adults. *Circulation* 2004;109(17):2068-2073.
15. Ailani J, Burch RC, Robbins MS et al. The American Headache Society consensus statement: update on integrating new migraine treatments into clinical practice. *Headache* 2021;61:1021-1039.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created from CP.PMN.122 for HFS migration	06.26.23	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted

standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2007 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.