

# Quick Reference Guide HEDIS® MY 2022



## Representation For more information, visit www.ncqa.org

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA). The HEDIS® measures and specifications were developed by and are owned by NCQA. NCQA holds a copyright in these materials and may rescind or alter these materials at any time. Users of the HEDIS® measures and specifications shall not have the right to alter, enhance, or otherwise modify the HEDIS® measures and specifications, and shall not disassemble, recompile, or reverse engineer the HEDIS® measures and specifications. Anyone desiring to use or reproduce the materials, subject to licensed user restrictions, without modification for an internal non-commercial purpose may do so without obtaining any approval from NCQA. Use of the Rules for Allowable Adjustments of HEDIS® to make permitted adjustments of the materials does not constitute a modification. All other uses, including a commercial use, or any external reproduction, distribution, or publication must be approved by NCQA and are subject to a license at the discretion of NCQA.

## HEDIS® MY 2022 Quick Reference Guide

Updated to reflect NCQA HEDIS® MY 2022 Technical Specifications

YouthCare strives to provide quality healthcare to our membership as measured through HEDIS® quality metrics. We created the HEDIS® MY 2022 Quick Reference Guide to help you increase your practice's HEDIS® rates and to use to address care opportunities for your patients. Please always follow the state and/ or CMS billing guidance and ensure the HEDIS® codes are covered prior to submission.

## What is HEDIS®?

HEDIS® (Healthcare Effectiveness Data and Information Set) is a set of standardized performance measures developed by the National Committee for Quality Assurance (NCQA) to objectively measure, report, and compare quality across health plans. NCQA develops HEDIS® measures through a committee represented by purchasers, consumers, health plans, health care providers, and policy makers.

## What Are the Scores Used For?

As state and federal governments move toward a quality-driven healthcare industry, HEDIS® rates are becoming more important for both health plans and individual providers. State purchasers of healthcare use aggregated HEDIS® rates to evaluate health insurance companies' efforts to improve preventive health outreach for patients.

Physician-specific scores are also used to measure your practice's preventive care efforts. Your practice's HEDIS® score



determines your rates for physician incentive programs that pay you an increased premium — for example Pay For Performance or Quality Bonus Funds.

## How Are Rates Calculated?

HEDIS® rates can be calculated in two ways: administrative data or hybrid data. Administrative data consists of claim or encounter data submitted to the health plan. Hybrid data consists of both administrative data and a sample of medical record data. Hybrid data requires review of a random sample of member medical records to abstract data for services rendered but that were not reported to the health plan through claims/encounter data. Accurate and timely claim/encounter data reduces the need for medical record review. If services are not billed or not billed accurately, they are not included in the calculation.

## **How Can I Improve My HEDIS® Scores?**

- Submit claim/encounter data for each and every service rendered
- Make sure that chart documentation reflects all services billed
- Bill (or report by encounter submission) for all delivered services, regardless of contract status
- Ensure that all claim/encounter data is submitted in an accurate and timely manner
- Consider including CPT II codes to provide additional details and reduce medical record requests

## Questions?



ILYouthCare.com



844-289-2264

Providers and other health care staff should document to the highest specificity to aid with the most correct coding choice.

Ancillary staff: Please check the tabular list for the most specific ICD-10 code choice.

This guide has been updated with information from the release of the HEDIS® MY 2022 Volume 2 Technical Specifications by NCQA and is subject to change.



For more information, visit www.ncqa.org

## **Contents**

#### Partnering with YouthCare 5 Introduction Consumer Assessment of Healthcare Providers and Systems (CAHPS®) 10 **Cultural Competence** Quality Provider Webinar (QPW) 11 12 Access & Availability 15 Caring for and Communicating with Individuals with Intellectual and Developmental Disabilities (IDD) 18 **Adult Health** 18 Adults' Access to Preventive/Ambulatory Health Services (AAP) 19 Antidepressant Medication Management (AMM) 20 Controlling High Blood Pressure (CBP) 21 Blood Pressure Control for Patients with Diabetes (BPD)\* 21 Eye Exam for Patients with Diabetes (EED)\* 22 Hemoglobin A1c Control for Patients with Diabetes (HBD)\* 23 Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD) 24 Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) 25 Women's Health 25 Chlamydia screening in Women (CHL) 26 Prenatal and Postpartum Care (PPC) 27 **Pediatric Health** 27 Follow-Up Care for Children Prescribed ADHD Medication (ADD) 29 Annual Dental Visit (ADV) 30 Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM) 31 Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) 32 Childhood Immunization Status (CIS) 33 Immunizations for Adolescents (IMA) 34 Lead Screening in Children (LSC) Well-Child and Adolescent Well-Care Visits (W30/WCV) 35 36 Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) **37** General Health 37 Asthma Medication Ratio (AMR) 39 Appropriate Testing for Pharyngitis (CWP)

- 40 Follow-Up After Emergency Department Visit for Substance Use (FUA)
- 41 Follow-Up After Hospitalization for Mental Illness (FUH)
- 43 Follow-Up After Emergency Department Visit for Mental Illness (FUM)
- 44 Initiation and Engagement of Substance Use Disorder Treatment (IET)

#### 47 Appendix

- 47 Point of Service Code Definitions
- 49 Exclusions

# Partnering with YouthCare

## Introduction

This guide contains information about the quality measures for the YouthCare

# Consumer Assessment of Healthcare Providers and Systems (CAHPS®)

Every year, a random sample of YouthCare patients are surveyed about their experience with their providers, services, and health plan. It is an important component of ensuring that patients are satisfied, not only with their health outcomes but also with their healthcare experience.

CAHPS® surveys allow patients to evaluate the aspects of care delivery that matter the most to them. At YouthCare, we are committed to partnering with our providers to deliver an outstanding patient experience.

As a provider, you are the most critical component of that experience. We want to ensure that you know exactly how your patients are evaluating your care. Please take a moment to review and to familiarize yourself with some of the key topics included in the survey.

CAHPS® Measure	Description	Daily Practice Tips
Getting Needed Care	This measure assesses the ease with which patients received the care, tests, or treatment they needed. It also assesses how often they were able to get a specialist appointment scheduled when needed.	Office staff should help coordinate specialty appointments for urgent cases Encourage patients and caregivers to view results on the patient portal when available Inform patients of what to do if care is needed after hours Offer appointments or refills via text and/or email Offer alternative appointment types to expand access to care (e.g., telephone, telehealth, telemedicine, and patient portals)
Getting Care Quickly	This measure assesses how often patients got the care they needed as soon as they needed it and how often appointment wait times exceeded 15 minutes.	Ensure a few appointments each day are available to accommodate urgent visits     Offer appointments with a nurse practitioner or physician assistant for short notice appointments     Maintain an effective triage system to ensure that frail and/or very sick patients are seen right away or provided alternate care via phone and urgent care     Keep patients informed if there is a longer wait time than expected and give them an option to reschedule

CAHPS® Measure	Description	Daily Practice Tips
Care Coordination	This measure assesses providers' assistance with managing the disparate and confusing health care system, including access to medical records, timely follow-up on test results, and education on prescription medications.	<ul> <li>Ensure there are open appointments for patients recently discharged from a facility</li> <li>Integrate PCP and specialty practices through EMR or fax to get reports promptly</li> <li>Ask patients if they have seen any other providers; discuss visits to specialty care as needed</li> <li>Encourage patients to bring in their medications to each visit</li> </ul>
How Well Doctors Communicate	This measure assesses patients' perception of the quality of communication with their doctor. Consider using the Teach-Back Method to ensure patients understand their health information.	What is Teach-back?  • A way to ensure you – the healthcare provider – have explained information clearly. It is not a test or quiz of patients  • Asking a patient (or family member) to explain in their own words what they need to know or do, in a caring way  • A way to check for understanding and, if needed, re-explain and check again  • A research-based health literacy intervention that improves patient-provider communication and patient health outcomes

CAHPS® Measure	Description	Daily Practice Tips
Rating of Health Care Quality	The CAHPS® survey asks patients to rate the overall quality of their health care on a 0-10 scale.	Encourage patients to make their routine appointments for checkups or follow up visits as soon as they can – weeks or even months in advance     Ensure that open care gaps are addressed during each patient visit     Make use of the provider portal when requesting prior authorizations

On the following page are examples of satisfaction categories and survey questions for which your patients are asked to respond; provider discussion questions; and Provider Tips. We hope this tool will provide reinforcement opportunities for your relationship with the patients you serve.

Sample Questions	Provider Tips	
Health Promotion Discussion Questions Any problems with your work or daily	Complete and document any     health assessment on nations.	
activities due to physical problems?  Any problems with your work or daily activities due to stress?  Anything bothering you or stressful?	<ul> <li>health assessment on patient</li> <li>Discuss with patient the benefits of exercise and encourage them to start, increase or maintain physical</li> </ul>	
Are you sad or depressed?	activity and document discussion	
Do you use tobacco? (Always/Sometimes/Never)	Discuss the risks of tobacco use and recommend medication to assist in stopping	
Do you drink alcohol? (Always/Sometimes/Never)	Discuss issues associated with drinking too much alcohol,	
Do you exercise? (Always/Sometimes/Never)	if necessary	
Do you take aspirin? (Always/Sometimes/Never)	Discuss the risks and benefits of aspirin to prevent heart attack or stroke	
Do you or anyone in your family have high blood pressure, high cholesterol or had a heart attack?	Screen patient for high blood pressure and cholesterol	
Have you had a flu shot in the past calendar year? If not, Why?	Recommend and/or administer the flu shot during flu season	

Sample Questions	Provider Tips
Medication Discussion Questions	
Are you currently on any prescription medications from another doctor? If so, what?	Document all prescription medication patient is taking
How long have you been on the medication?	<ul> <li>Discuss options and reasons to take alternate medications if patient is not getting positive results for symptoms</li> <li>Discuss reasons with patient why they may need to stop taking a particular medication</li> <li>Discuss the benefits and risks of taking a medicine</li> <li>Discuss patient's preference on what medication they feel would be best for them</li> <li>Review medications prescribed by PCP and specialists and verify results</li> </ul>
Access to Care Discussion Questions	
Are you satisfied with the timeframe it took to schedule your appointment?	Evaluate office procedures to improve getting patients scheduled as quickly as possible for their symptoms
Were you able to get your appointment as soon as you needed?	Determine why patient perceives difficulty in getting timely care, if necessary
Are you satisfied with the coordination of care you receive, coordinating visits with specialists, non-emergency transportation (if needed) and providing lab or test results?	<ul> <li>Educate patient on timeframes for getting appointments according to their symptoms</li> <li>Assist in coordination of non-emergency transportation, if necessary</li> </ul>

## **Cultural Competence**

Cultural Competence is a set of attitudes, behaviors, and policies that enable people to work effectively in cross-cultural situations. We serve a diverse patient population. The ability to understand and relate to different cultures can help you communicate effectively with your patients. All YouthCare network providers are contractually required to complete the on-line Cultural Competence training module annually.

## Tips for Providing Culturally Competent Care

Consider population-specific conditions: Low-income/ low-literacy, race, disability, spirituality, age, sexuality, and gender identity.

Ask about cultural practices: Spiritual traditions, dietary restrictions, and more may impact a patient's clinical experience.

Practice transcultural techniques: Approach a new patient slowly, be respectful, sit in a quiet setting, and sit a comfortable distance away.

Ensure patient's understanding of care: Lack of accessible medication instructions in a patient's language can impact quality of care. Ensure a patient's comprehension by utilizing translated handouts and/or make use of a translator.

## Things to Remember

- 1 in 4 Americans live with a disability and are twice as likely to find his or her provider's skills or facilities inadequate.
- 1 in 5 Americans speak a language other than English at home. Language barriers can prevent patients from effectively conveying their ailments and understanding their care plans.
- 3.6 million Americans miss or delay medical care because they lack reliable transportation.
- Invest in Americans with Disabilities Act (ADA)-approved renovations and train staff on disabilities, challenges, and rights.
- Speak slowly, summarize, demonstrate, and use appropriate terminology when providing instructions. Ensure that patients understand the instructions at the end of the visit. Use PALS, YouthCare's interpreter service, to better serve YouthCare patients. To reach PALS, please call 313-351-9388 and provide the customer code 7119.
- Call YouthCare transportation at 866-796-1165 at least three business days prior to a patient's appointment.

Sources: CDC.gov, census.gov, ncbi.nlm.nih.gov

## **Quality Provider Webinar (QPW)**

To support our providers in their quality improvement efforts, YouthCare's Quality Improvement team hosts a monthly one hour webinar on topics related to improving patients' quality of care including a monthly presentation that spotlights a clinical area of focus.

We hope these sessions will assist provider teams to improve HEDIS® scores and drive better incentive payments to your practice.

All office staff can attend including providers, administrative staff, and quality teams. Participants can watch the webinars remotely using the Zoom call-in number sent each month via email. If you are interested in attending a session or receiving a copy of a webinar presentation or 2022 webinar schedule, email illinoisdataabstractionteam@mhplan.com.

The webinars take place on the third Thursday of every month. Webinars are currently scheduled as live sessions from 12:00 p.m. CT to 1:00 p.m. CT.

#### Sample webinar topics include:

- HEDIS® Exclusions and Medical Record Retrieval Methods
- Adult and Behavioral Health HEDIS® Measures
- Children's HFDIS® Measures
- Patient Satisfaction, Health Outcomes Survey and Patient-Centered Medical Homes

## **Access & Availability**

Annually, YouthCare assesses the appointment availability and after-hours access of its contracted provider offices to ensure patients are served based on their level of need

Each year, our Quality Improvement team conducts a telephone audit using the standards below set forth by NCQA, CMS and/or State regulations. These audits are conducted in an effort to monitor provider compliance with Illinois Medicaid contract requirements. The process of conducting the annual audits is outlined below.

IDENTIFY YouthCare audits a sample of its contracted PCPs, behavioral health practitioners and specialists.

OUTREACH Conducted by the Quality Improvement team via phone, three attempts are made to reach a live person.

ANALYZE Analysis is performed based on all data collected.

**REPORT** Letters are mailed to offices indicating the results of the audit. Any offices who did not meet the standards are placed on a corrective action plan and are asked to identify ways to improve their appointment availability.

### Appointment Availability Standards

Appointment Types	Population	Standard
Medicaid	T opalation	Starradia
Preventive/Routine Care	Child ≤ 6 Months	2 weeks
Preventive/Routine Care	Child > 6 Months	5 weeks
Preventive/Routine Care	Adult	5 weeks
Urgent/Non-Emergent (Medically Necessary Care)	Adult or Child	1 business day (24 hours)
Non-Urgent/Non-Emergent Conditions	Adult or Child	3 weeks
Initial Prenatal w/o Problems (First Trimester)	Female Enrollees	2 weeks
Prenatal (Second Trimester)	Female Enrollees	1 week
Prenatal (Third Trimester)	Female Enrollees	3 days
Office Wait Time	All	Less than 30 minutes
Hours Different for Medicaid	All	No; must
Recipients		be same
Medicare		
Preventive/Routine Care	Adult	5 weeks

Appointment Types	Population	Standard
Non-Urgent/Non-Emergent Conditions	Adult	3 weeks
Urgent/Non-Emergent (Medically Necessary Care)	Adult	24 hours
Emergency	Adult	Immediate after request or referred to ED
Initial Prenatal w/o Problems (First Trimester)	Female Enrollees	2 weeks
Prenatal (Second Trimester)	Female Enrollees	1 week
Prenatal (Third Trimester)	Female Enrollees	3 days
Office Wait Time	All	Less than 30 minutes

Medical coverage 24 hours a day, 7 days a week

## Behavioral Health Appointment Availability Standards

Appointment Types	Standard
Life Threatening Emergency	Immediately, or referred to the Emergency Room
Initial Visit for Routine Care	Within 10 business days
Follow-Up Visit for Routine Care	Within 14 business days
Office Wait Time	< 30 minutes
Scheduled Appointments per Hour, per Practitioner	≤ 6 per hour
Different Hours for Medicaid	No; must be the same

Medical coverage 24 hours a day, 7 days a week

## Specialist Appointment Availability Standards

Appointment Types	Standard	Population
Medicaid		
Routine Office Visit	Adult	Within 45 calendar days
Routine Office Visit	Child	Within 21 calendar days
Urgent Visit	All	Within 72 hours
Office Wait Time	All	< 30 minutes
Scheduled Appointments per Hour, per Practitioner	All	≤ 6 per hour
Hours Different for Medicaid and Medicare Recipients	All	No; must be the same

#### Appointment Availability Standards (continued)

Medical coverage 24 hours a day, 7 days a week

#### After-hours standards

All specialist contracts require practitioners to ensure coverage for their respective practices 24 hours a day, seven days a week.

#### Acceptable after-hours access mechanisms include:

- Answering service
- On-call pager/cellular
- Call forwarded to practitioner's home or other location
- Published after-hours telephone number and recorded voice message directing patients to a practitioner for urgent and non-life threatening conditions. The message should not instruct patients to obtain treatment at the Emergency Room for non-life-threatening emergencies.

#### Message Components

• Message MUST direct patients in a medical emergency to call 911 or go to the nearest Emergency Room or Urgent Care.

#### Message Must Contain ONE of the Following:

- Message forwards to on-call practitioner
- Message forwards to an answering service
- Message gives the on-call practitioner's number
- Message gives the on-call practitioner's pager
- · Message refers patient to another office, practitioner, or on-call service Message may not only direct patient to Emergency Room. The patient must be able to leave a message for an on-call doctor, speak with an on-call doctor or be forwarded to an on-call doctor.

## Caring for and Communicating with Individuals with Intellectual and Developmental Disabilities (IDD)

As a provider, it is important to be aware of the following health disparities individuals with intellectual and developmental disabilities may face:

- Fewer preventive screenings than the general population
- · Financial and transportation limitations
- Lack of access to specialized training or experience with caring for individuals in these populations

## Complete an Annual Functional Status Assessment:

This assessment measures the patient's ability to perform daily tasks and helps to identify any functional decline. For YouthCare patients, please indicate one of the following in the medical record:

- Notation and date that Activities of Daily Living (ADL) were assessed -Bathing, dressing, eating, transferring, using toilet, walking
- Notation and date that Instrumental Activities of Daily Living (IADL) were assessed - Shopping, driving or using public transportation, meal preparation, housework, taking medications, using the telephone
- Result of a standardized functional status assessment and the date it was performed - Assessment of Living Skills and Resources (ALSAR), Barthel ADL Index Physical Self-Maintenance (ADLS) Scale, Bayer Activities of Daily Living (B-ADL) Scale, Extended Activities of Daily Living (EADL) Scale
- · Chronic conditions at a younger age
- A higher risk of obesity
- · A greater incidence of mental illness
- An accelerated aging process

It is important that providers and health plans adapt and coordinate care for this population by communicating appropriately and respectfully about individuals with disabilities.

Individuals with intellectual and developmental disabilities often go through cognitive and behavioral changes. These include anxiety, depression, dementia, self-injurious behavior, and other factors. These changes may result from pain and discomfort related to other medical issues.

As a provider, it is important to be aware of the following ways to communicate, in order to emphasize the person first, and not the disability:

- Emphasize abilities, not limitations
- Do not use language that suggests the lack of something
- Emphasize the need for accessibility, not the disability
- Do not use offensive language
- Avoid language that implies negative stereotypes
- Do not portray people with disabilities as inspirational only because of their disability

## Tips on Using People-First Language

This chart provides tips on what language to use when communicating with an individual with a disability.

Tips	Use	Do Not Use
Emphasize abilities, not limitations	Person who uses a wheelchair	Confined or restricted to a wheelchair, wheelchair bound
	Person who uses a device to speak	Can't talk, mute
Do not use	Person with a disability	Disabled, handicapped
language that	Person of short stature	Midget
suggests the lack of something	Person with cerebral palsy	Cerebral palsy victim
or something	Person with epilepsy or seizure disorder	Epileptic
	Person with multiple sclerosis	Afflicted by multiple sclerosis
Emphasize the need for accessibility, not the disability	Accessible parking or bathroom	Handicapped parking or bathroom
Do not use offensive language	Person with a physical disability	Crippled, lame, deformed, invalid, spastic
	Person with an intellectual, cognitive, developmental disability	Slow, simple, moronic, defective, afflicted, special person
	Person with an emotional or behavioral disability, a mental health impairment, or a psychiatric disability	Insane, crazy, psycho, maniac, nuts
Avoid language that implies negative stereotypes	Person without a disability	Normal person, healthy person
Do not portray people with disabilities as inspirational only because of their disability	Person who is successful, productive	Has overcome his/her disability, is courageous

Source: https://www.cdc.gov/ncbddd/disabilityandhealth/pdf/communicating-withpeople.pdf


#### Resources for Patients and Providers:

Illinois Department of Healthcare and Family Services (HFS): www.illinois.gov/hfs/MedicalClients/HCBS/Pages/support\_cyadd.aspx

Illinois Department of Human Services (IDHS): 217-782-3075

For a copy of current clinical practice guidelines, visit our website at ILYouthCare.com.

Provider Resources: <a href="https://www.ilyouthcare.com/providers/provider-">https://www.ilyouthcare.com/providers/provider-</a> resources.html



# **Adult** Health

## **Adults' Access to Preventive/Ambulatory Health Services (AAP)**

The AAP measure evaluates the percentage of patients 20 years and older who had an ambulatory or preventive care visit. Services that count include outpatient evaluation and management (E&M) visits, consultations, assisted living/home care oversight, preventive medicine, and counseling.

#### **AAP Measure Codes**

CPT*	HCPCS*	ICD-10*
92002, 92004, 92012, 92014,	G0071, G0402, G0438,	Z00.00, Z00.01,
98966-98972, 99201-99205,	G0439, G0463, G2010,	Z00.3, Z00.5,
99211-99215, 99241-99245,	G2012, G2061, G2062,	Z00.8, Z00.121,
99304-99310, 99315, 99316,	G2063, T1015, S0620,	Z00.129, Z02.0,
99318, 99324-99328, 99334-	S0621	Z02.1, Z02.2,
99337, 99341-99345, 99347-		Z02.3, Z02.4,
99350, 99381-99387, 99391-		Z02.5, Z02.6,
99397, 99401-99404, 99411,		Z02.71, Z02.79,
99412, 99421-99423, 99429,		Z02.81, Z02.82,
99441-99443, 99444, 99457,		Z02.83, Z02.89,
99483		Z02.9, Z76.1, Z76.2

<sup>\*</sup>Codes subject to change

## **Antidepressant Medication Management (AMM)**

The AMM measure evaluates the percentage of patients 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an antidepressant medication treatment.

#### Two rates are reported:

- Effective Acute Phase Treatment: percentage of patients who remained on an antidepressant medication for at least 84 days (12 weeks)
- Effective Continuation Phase Treatment: percentage of patients who remained on an antidepressant medication for at least 180 days (6 months)

## **Antidepressant Medications**

Description	Prescription	
Miscellaneous antidepressants	<ul><li>Bupropion</li><li>Vortioxetine</li></ul>	• Vilazodone
Monoamine oxidase inhibitors	<ul><li>Isocarboxazid</li><li>Phenelzine</li></ul>	<ul><li>Selegiline</li><li>Tranylcypromine</li></ul>
Phenylpiperazine antidepressants	• Nefazodone	• Trazodone
Psychotherapeutic combinations	<ul><li>Amitriptyline-chlordiazepoxide</li><li>Fluoxetine-olanzapine</li><li>Amitriptyline-perphenazine</li></ul>	
SNRI antidepressants	<ul><li>Desvenlafaxine</li><li>Venlafaxine</li></ul>	<ul><li>Duloxetine</li><li>Levomilnacipran</li></ul>
SSRI antidepressants	<ul><li>Citalopram</li><li>Fluoxetine</li><li>Paroxetine</li></ul>	<ul><li>Escitalopram</li><li>Fluvoxamine</li><li>Sertraline</li></ul>
Tetracyclic antidepressants	Maprotiline	Mirtazapine
Tricyclic antidepressants	<ul><li>Amitriptyline</li><li>Clomipramine</li><li>Doxepin (&gt;6 mg)</li><li>Nortriptyline</li><li>Trimipramine</li></ul>	<ul><li>Amoxapine</li><li>Desipramine</li><li>Imipramine</li><li>Protriptyline</li></ul>

## **Controlling High Blood Pressure (CBP)**

The CBP measure evaluates the percentage of patients 18-85 years of age who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled (<140/90 mm Hg).

All recorded blood pressure measurements

- Consider taking two readings at each office visit. Sometimes a second blood pressure reading is lower
- Do not round blood pressure; always use exact numbers

#### **CBP Measure Codes**

Description	Codes*
Essential Hypertension	ICD-10: I10
Systolic Greater Than/Equal to 140	CPT-CAT-II: 3077F
Systolic 130-139	CPT-CAT-II: 3075F
Systolic Less Than 130	CPT-CAT-II: 3074F
Diastolic Greater Than/Equal to 90	CPT-CAT-II: 3080F
Diastolic 80-89	CPT-CAT-II: 3079F
Diastolic Less Than 80	CPT-CAT-II: 3078F

<sup>\*</sup>Codes subject to change

Comprehensive Diabetes Care (CDC) has been replaced by three separate indicators: Blood Pressure Control for Patients with Diabetes (BPD), Eye Exam for Patients with Diabetes (EED), and Hemoglobin A1c Control for Patients with Diabetes (HBD).

## **Blood Pressure Control for Patients with Diabetes (BPD)**

The BPD measure evaluates the percentage of patients 18-75 years of age with diabetes (types 1 and 2) whose blood pressure (BP) was adequately controlled (<140/90 mm Hg) during the measurement year.

#### **BPD Measure Codes**

Description	Codes*
Diastolic 80-89	CPT-CAT-II: 3079F
Diastolic Greater Than or Equal to 90	CPT-CAT-II: 3080F
Diastolic Less Than 80	CPT-CAT-II: 3078F
Systolic Greater Than or Equal to 140	CPT-CAT-II: 3077F
Systolic Less Than 130	CPT-CAT-II: 3074F
Systolic 130-139	CPT-CAT-II: 3075F

<sup>\*</sup>Codes subject to change

## **Eye Exam for Patients with Diabetes (EED)**

The EED measure evaluates percentage of patients 18-75 years of age with diabetes (types 1 and 2) who had a retinal eye exam.

#### **EED Measure Codes**

Description	Codes*
Unilateral Eye Enucleation with a Bilateral Modifier 50	<b>CPT</b> : 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114
Automated Eye Exam	CPT: 92229
Diabetic Retinal Screening Negative in Prior Year	CPT-CAT-II: 3072F
Eye Exam With Retinopathy	CPT-CAT-II: 2022F, 2024F, 2026F
Eye Exam Without Retinopathy	CPT-CAT-II: 2023F, 2025F, 2033F
Diabetes Mellitus Without Complications	ICD-10: E10.9, E11.9, E13.9

Description	Codes*
Diabetic Retinal Screening	CPT: 67028, 67030, 67031, 67036, 67039, 67040-67043, 67101, 67105, 67107, 67108, 67110, 67113, 67121, 67141, 67145, 67208, 67210, 67218, 67220, 67221, 67227, 67228, 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92225-92228, 92230, 92235, 92240, 92250, 92260, 99203-99205, 99213-99215,
	99242-99245 HCPCS: S0620, S0621, S3000

<sup>\*</sup>Codes subject to change

## **Hemoglobin A1c Control for Patients with Diabetes (HBD)**

The HBD measure evaluates the percentage of patients 18-75 years of age with diabetes (types 1 and 2) whose HbA1c was at the following levels: HbA1c control (<8.0%) and HbA1c poor control (>9.0%).

#### **HBD** Measure Codes

Description	Codes*
HbA1c Lab Test	CPT: 83036, 83037
HbA1c Level Greater Than 9.0	CPT-CAT-II: 3046F
HbA1c Level Greater Than or Equal to 7.0 and Less Than 8.0	CPT-CAT-II: 3051F
HbA1c Level Greater Than or Equal to 8.0 and Less Than or Equal to 9.0	CPT-CAT-II: 3052F
HbA1c Level Less Than 7.0	CPT-CAT-II: 3044F

<sup>\*</sup>Codes subject to change

## **Diabetes Monitoring for People with Diabetes and** Schizophrenia (SMD)

The SMD measure evaluates the percentage of patients 18-64 years of age with schizophrenia or schizoaffective disorder and diabetes who had both an LDL-C test and an HbA1c test.

### **SMD Measure Codes**

Description	Codes*
HbA1C Lab Tests	CPT: 83036, 83037
Most recent hemoglobin A1c (HbA1c) level less than 7.0% (DM)	CPT-CAT-II: 3044F
Most recent hemoglobin A1c level greater than 9.0% (DM)	CPT-CAT-II: 3046F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 7.0% and less than 8.0% (DM)	CPT-CAT-II: 3051F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 8.0% and less than or equal to 9.0% (DM)	CPT-CAT-II: 3052F

LDL-C Lab Tests	CPT: 80061, 83700, 83701, 83704, 83721
Most recent LDL-C less than 100 mg/dL (CAD) (DM)	CPT-CAT-II: 3048F
Most recent LDL-C 100-129 mg/ dL (CAD) (DM)	CPT-CAT-II: 3049F
Most recent LDL-C greater than or equal to 130 mg/dL (CAD) (DM)	CPT-CAT-II: 3050F

<sup>\*</sup>Codes subject to change

## **Diabetes Screening for People with Schizophrenia** or Bipolar Disorder Who Are Using Antipsychotic **Medications (SSD)**

The SSD measure evaluates percentage of patients 18-64 years of age with schizophrenia, schizoaffective disorder, or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test.

### SSD Measure Codes

Description	Codes*
HbA1C Lab Tests	CPT: 83036, 83037
Most recent hemoglobin A1c (HbA1c) level less than 7.0% (DM)	CPT-CAT-II: 3044F
Most recent hemoglobin A1c level greater than 9.0% (DM)	CPT-CAT-II: 3046F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 7.0% and less than 8.0% (DM)	CPT-CAT-II: 3051F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 8.0% and less than or equal to 9.0% (DM)	CPT-CAT-II: 3052F
Glucose Lab Tests	CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951

<sup>\*</sup>Codes subject to change

# Women's Health

## **Chlamydia screening in Women (CHL)**

The CHL measure evaluates the percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia.

**CHL Measure Codes** 

CPT\*

87110, 87270, 87320, 87490-87492, 87810

\*Codes subject to change

## **Prenatal and Postpartum Care (PPC)**

The PPC measure evaluates percentage of deliveries of live births on or between October 8 of the year prior to the measurement year and October 7 of the measurement year. For these women, the measure assesses the following facets of prenatal and postpartum care:

- Timeliness of Prenatal Care: percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date, or within 42 days of enrollment in the organization
- Postpartum Care: percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery

#### **PPC Measure Codes**

Description	Codes*
Online Assessments	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061, G2062, G2063
Prenatal Visits	CPT: 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99483 HCPCS: G0463, T1015
Stand-Alone Prenatal Visits	CPT: 99500 CPT-CAT-II: 0500F, 0501F, 0502F HCPCS: H1000, H1001, H1002, H1003, H1004
Prenatal Bundled Services	CPT: 59400, 59425, 59426, 59510, 59610, 59610, 59618 HCPCS: H1005
Cervical Cytology Lab Test	CPT: 88141-88143, 88147, 88148, 88150, 88152-88153, 88164-88167, 88174, 88175 HCPCS: G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091
Postpartum Visits	CPT: 57170, 58300, 59430, 99501 CPT-CAT-II: 0503F HCPCS: G0101 ICD-10: Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2
Telephone Visits	CPT: 98966-98968, 99441-99443
Postpartum Bundled Services	CPT: 59400, 59410, 59510, 59515, 59610, 59614, 59618, 59622

<sup>\*</sup>Codes subject to change

NOTE: When using the Online Assessment, Telephone Visit, or Prenatal Visit codes, remember to also include a Pregnancy Diagnosis code.



# **Pediatric** Health

## Follow-Up Care for Children Prescribed ADHD Medication (ADD)

The ADD measure evaluates percentage of children newly prescribed attention deficit hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed.

## Two rates are reported:

- Initiation Phase: percentage of patients 6-12 years of age with a prescription dispensed for ADHD medication, who had one follow-up visit with a practitioner with prescribing authority during the 30-day Initiation Phase
- Continuation and Maintenance (C&M) Phase: percentage of patients 6-12 years of age with a prescription dispensed for ADHD medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended.

## **ADD Measure Codes**

Description	Codes*
An Outpatient Visit	CPT: 90791, 90792, 90832-90834, 90836- 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
BH Outpatient Visit	CPT: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99510, 99483 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015
Observation Visit	CPT: 99217-99220
Health and Behavior Assessment/Intervention	CPT: 96150-96154
Visit Setting Unspecified Value Set with Partial Hospitalization POS	CPT: 90791, 90792, 90832-90834, 90836- 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 52
Partial Hospitalization/ Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Visit Setting Unspecified Value Set with Community Mental Health Center POS	CPT: 90791, 90792, 90832-90834, 90836- 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 53

<sup>\*</sup>Codes subject to change

#### **ADHD Medications**

CNS Stimulants		
<ul><li>Dexmethylphenidate</li><li>Dextroamphetamine</li><li>Lisdexamfetamine</li></ul>	Methamphetamine     Methylphenidate	
Alpha-2 Receptor Agonists:		
Clonidine	Guanfacine	
Misc. ADHD Medication:		
Atomoxetine		

## **Annual Dental Visit (ADV)**

The ADV measure evaluates the percentage of patients two to 20 years of age who had at least one dental visit during the measurement year.

YouthCare partners with Envolve Dental to administer dental benefits for patients. Envolve Dental can be reached at 855-586-1415.

Primary Care Providers (PCPs) can receive training to administer oral health screenings and fluoride varnish and sealants by completing the Children's Oral Health Smiles for Life Course 6: Caries Risk Assessment, Fluoride Varnish, and Counseling training module. By completing this course, PCPs can help prevent their patients from developing serious health conditions.

The training module can be found at www.smilesforlifeoralhealth.org.

## **Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)**

The APM measure evaluates the percentage of children and adolescents 1-17 years of age who had two or more antipsychotic prescriptions and had metabolic testing.

### Three rates are reported:

- Percentage of children and adolescents on antipsychotics who received blood glucose testing
- Percentage of children and adolescents on antipsychotics who received cholesterol testing
- Percentage of children and adolescents on antipsychotics who received blood glucose and cholesterol testing

#### **APM Measure Codes**

Description (Need either A1c or Glucose and LCL-C or Cholesterol)	Codes*
HbA1C Lab Tests	ICD-10: 83036, 83037
HbA1c Level Greater Than 9.0	CPT-CAT-II: 3046F
HbA1c Level Greater Than or Equal To 7.0 and Less Than 8.0	CPT-CAT-II: 3051F
HbA1c Level Greater Than or Equal To 8.0 and Less Than or Equal To 9.0	CPT-CAT-II: 3052F
HbA1c Level Less Than 7.0	CPT -CAT-II: 3044F

Glucose Lab Tests	CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951
LDL-C Lab Tests	CPT: 80061, 83700, 83701, 83704, 83721
Most recent LDL-C less than 100 mg/dL (CAD) (DM)	CPT-CAT-II: 3048F
Most recent LDL-C 100-129 mg/dL (CAD) (DM)	CPT-CAT-II: 3049F
Most recent LDL-C greater than or equal to 130 mg/dL (CAD) (DM)	CPT-CAT-II: 3050F

<sup>\*</sup>Codes subject to change

## **Use of First-Line Psychosocial Care for Children** and Adolescents on Antipsychotics (APP)

The APP measure evaluates the percentage of children and adolescents 1 to 17 years of age who had a new prescription for an antipsychotic medication and had documentation of psychosocial care as first line treatment.

## **Antipsychotic Medications**

Miscellaneous Antipsychotic Agents		
<ul> <li>Aripiprazole</li> <li>Asenapine</li> <li>Brexpiprazole</li> <li>Cariprazine</li> <li>Clozapine</li> <li>Haloperidol</li> </ul>	<ul><li>Iloperidone</li><li>Loxapine</li><li>Lurisadone</li><li>Molindone</li><li>Olanzapine</li><li>Paliperidon</li></ul>	<ul><li>Pimozide</li><li>Quetiapine</li><li>Risperidone</li><li>Ziprasidone</li></ul>
Phenothiazine Antipsychotics		
<ul><li>Chlorpromazine</li><li>Fluphenazine</li></ul>	<ul><li>Perphenazine</li><li>Thioridazine</li></ul>	Trifluoperazine
Thioxanthenes		
Thiothixene		
Long-Acting Injections		
<ul><li> Arirprazole</li><li> Arirprazole lauroxil</li></ul>	<ul><li> Haloperidol decanoate</li><li> Olanzapine</li></ul>	Risperidone

• Paliperidone palmitate

#### **Antipsychotic Combination Medications**

#### **Psychotherapeutic Combinations**

- Fluoxetine-olanzapine
- Perphenazine-amitriptyline

#### APP Measure Codes

Fluphenazine

decanoate

Description	CPT*	HCPCS*
Psychosocial Care	90832-90834,	G0409, G0410, H0004,
	90836-90840,	H0035-H0040, H2000,
	90845-90847, 90849,	H20001, H2011-H2014,
	90853, 90875, 90876,	H2017-H2020, S0201,
	90880	S9480, S9484, S9485

<sup>\*</sup>Codes subject to change

## **Childhood Immunization Status (CIS)**

The CIS measure evaluates the percentage of children 2 years of age who completed immunizations on or before child's second birthday.

## CIS Measure Codes

Description	Codes*
DTaP (4 dose)	CPT: 90697, 90698, 90700, 90723 CVX: 20, 50, 106, 107, 110, 120, 146
HIB (3 dose)	CPT: 90644, 90647, 90648, 90697, 90698, 90748 CVX: 17, 46, 47, 48, 49, 50, 51, 120, 146, 148
Newborn Hep B (3 dose)	CPT: 90697, 90723, 90740, 90744, 90747, 90748 CVX: 08, 44, 45, 51, 110, 146 HCPCS: G0010 ICD-10-CM: B16.0, B16.1, B16.2, B16.9, B17.0, B18.0, B18.1, B19.10, B19.11, 3E0234Z
IPV (3 dose)	CPT: 90697, 90698, 90713, 90723 CVX: 10, 89, 110, 120, 146
MMR (1 dose)	CPT: 90707, 90710 CVX: 03, 94 ICD-10: B05.0, B05.1, B05.2, B05.3, B05.4, B05.81, B05.89, B05.9, B26.0, B26.1, B26.2, B26.3, B26.81, B26.82. B26.83, B26.84, B26.85, B26.89, B26.9, B06.00, B06.01, B06.02, B06.09, B06.81, B06.82, B06.89, B06.9
Pneumococcal Conjugate PCV (4 dose)	CPT: 90670 CVX: 109, 133, 152 HCPCS: G0009
Varicella VZV (1 dose)	CPT: 90710, 90716 CVX: 21, 94 ICD-10: B01.0, B01.11, B01.12, B01.2, B01.81, B01.89, B01.9, B02.0, B02.1, B02.21, B02.22, B02.23, B02.24, B02.29, B02.30, B02.31, B02.32, B02.33, B02.34, B02.39, B02.7, B02.8, B02.9
Hep A (1 dose)	CPT: 90633 CVX: 31, 83, 85 ICD-10: B15.0, B15.9
Influenza Flu (2 dose) LAIV vaccination must be administered on the child's 2nd birthday	CPT: 90655, 90657, 90660, 90661, 90672, 90673, 90674, 90685-90689, 90756 CVX: 88, 140, 141, 150, 153, 155, 158, 161, 111, 149, 171, 186 HCPCS: G0008

Description	Codes*
Rotavirus (2 Dose)	CPT: 90681 CVX: 119
Rotavirus (3 Dose)	CPT: 90680 CVX: 116, 122

<sup>\*</sup>Codes subject to change

NOTE: Rotavirus is either 2 dose OR 3 dose for compliancy

## **Immunizations for Adolescents (IMA)**

The IMA measure evaluates percentage of adolescents 13 years of age who completed immunizations on or before patient's 13th birthday.

## **IMA Measure Codes**

Description	Codes*
Meningococcal-serogroup A,C,W, and Y: (1 dose)	CPT: 90619, 90733, 90734 CVX: 32, 108, 114, 136, 147, 167, 203
Tdap (1 dose)	CPT: 90715 CVX: 115
HPV (2 or 3 dose series)	CPT: 90649-90651 CVX: 62, 118, 137, 165

<sup>\*</sup>Codes subject to change

## Lead Screening in Children (LSC)

The LSC measure evaluates percentage of children 2 years of age who had one or more capillary or venous lead blood test for lead poisoning on or before members' second birthday.

#### LSC Measure Code

CPT\*

83655

\*Codes subject to change

In order to ensure that children receive appropriate public health follow-up services, physicians and other health providers have an obligation to report blood lead results greater than or equal to 10mg/dL within 48 hours to the Illinois Department of Public Health (IDPH) Childhood Lead Poisoning Reporting System. Providers using the IDPH laboratory are not required to report blood lead results.

Illinois Department of Public Health Illinois Lead Program 535 W. Jefferson Street Springfield, IL 62761 P: 217-782-3517 www.idph.state.il.us

The Illinois Department of Healthcare and Family Services (HFS) encourages providers to send all blood lead specimens to the IDPH laboratory for analysis. Providers who utilize the state laboratory for blood lead analysis can order supplies for blood lead specimen collection free of charge by calling the IDPH Laboratory Shipping Section at 217-524-6222, or by downloading the Clinical Supplies Requisition Form from the HFS website: https://www.dph.illinois.gov/.

## Well-Child and Adolescent Well-Care Visits (W30/WCV)

The W30/WCV measure evaluates the percentage of patients within designated ages who had comprehensive well-care visit(s) as defined in each measure, with a PCP or an OB/GYN practitioner during the measurement year.

#### Well-Child Vists in the First 30 Months of Life (W30)

Months of Life: The percentage of patients who had the following number of well-child visits with a PCP during the last 15 months.

#### Two rates are reported:

- Well-Child Visits in the First 15 Months. Children who turned 15 months old during the measurement year: Six or more well-child visits.
- Well-Child Visits for Age 15 Months-30 Months. Children who turned 30 months old during the measurement year: Two or more well-child visits.

#### W30 Measure Codes

CPT*	HCPCS*	ICD-10*
99381-99385,	G0438, G0439, S0302	Z00.110, Z00.111, Z00.121,
99391-99395, 99461		Z00.129, Z00.2, Z76.1, Z76.2

<sup>\*</sup>Codes subject to change

(WCV) Child and Adolescent Well-Care Visits: Patients 3-21 years of age who had a least one comprehensive well-care visit with a PCP or an OB/GYN

#### **WCV Measure Codes**

CPT*	HCPCS*	ICD-10*
99381-99385,	G0438, G0439, S0302,	Z00.121, Z00.129, Z00.2,
99391-99395, 99461	S0610, S0612, S0613	Z00.3, Z01.411, Z01.419,
		Z76.1

<sup>\*</sup>Codes subject to change

## **Weight Assessment and Counseling for Nutrition** and Physical Activity for Children/Adolescents (WCC)

The WCC measure evaluates the percentage of patients 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of each of the following:

- BMI Percentile
- Counseling for Nutrition
- · Counseling for physical activity

### **WCC Measure Codes**

Description	Codes*
BMI Percentile	ICD-10: Z68.51, Z68.52, Z68.53, Z68.54
Nutrition Counseling	CPT: 97802-97804 HCPCS: G0270, G0271, G0447, S9449, S9452, S9470 ICD-10: Z71.3
Physical Activity	HCPCS: G0447, S9451 ICD-10: Z02.5, Z71.82

<sup>\*</sup>Codes subject to change



# General Health

## **Asthma Medication Ratio (AMR)**

The AMR measure evaluates the percentage of patients 5-64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medication of 0.50 or greater.

Oral medication dispensing event: One prescription of an amount lasting 30 days or less. Multiple prescriptions for different medications dispensed on the same day are counted as separate dispensing events.

Inhaler dispensing event: All inhalers of the same medication dispensed on the same day count as one dispensing event. Different inhaler medications dispensed on the same day are counted as different dispensing events.

**Injection dispensing event**: Each injection counts as one dispensing event. Multiple dispensed injections of the same or different medications count as separate dispensing events.

- Step 1: For each patient, count the units of asthma controller medications (Asthma Controller Medications List) dispensed during the measurement year.
- Step 2: For each patient, count the units of asthma reliever medications (Asthma Reliever Medications List) dispensed during the measurement year.
  - For each patient, sum the units calculated in step 1 and step 2 to determine units of total asthma medications
  - For each patient, calculate ratio using the below:
    - Units of Controller Medications/Units of Total Asthma Medications

## Asthma Controller Medications

Description	Prescriptions	Medication Lists	Route
Antibody inhibitors	Omalizumab	Omalizumab Medications List	Injection
Anti-interleukin-4	Dupilumab	Dupilumab Medications List	Injection
Anti-interleukin-5	Benralizumab	Benralizumab Medications List	Injection
Anti-interleukin-5	Mepolizumab	Mepolizumab Medications List	Injection
Anti-interleukin-5	Reslizumab	Reslizumab Medications List	Injection
Inhaled steroid combinations	Budesonide- formoterol	Budesonide Formoterol Medications List	Inhalation
Inhaled steroid combinations	Fluticasone- salmeterol	Fluticasone Salmeterol Medications List	Inhalation
Inhaled steroid combinations	Fluticasone- vilanterol	Fluticasone Vilanterol Medications List	Inhalation
Inhaled steroid combinations	Formoterol- mometasone	Formoterol Mometasone Medications List	Inhalation
Inhaled corticosteroids	Beclomethasone	Beclomethasone Medications List	Inhalation
Inhaled corticosteroids	Budesonide	Budesonide Medications List	Inhalation
Inhaled corticosteroids	• Ciclesonide	Ciclesonide Medications List	Inhalation
Inhaled corticosteroids	• Flunisolide	Flunisolide Medications List	Inhalation
Inhaled corticosteroids	• Fluticasone	Fluticasone Medications List	Inhalation
Inhaled corticosteroids	Mometasone	Mometasone Medications List	Inhalation
Leukotriene modifiers	Montelukast	Montelukast Medications List	Oral
Leukotriene modifiers	• Zafirlukast	Zafirlukast Medications List	Oral
Leukotriene modifiers	• Zileuton	Zileuton Medications List	Oral

## Asthma Controller Medications (continued)

Description	Prescriptions	Medication Lists	Route
Methylxanthines	Theophylline	Theophylline	Oral
		Medications List	

#### Asthma Reliever Medications

Description	Prescriptions	Medication Lists	Route
Short-acting, inhaled beta-2 agonists	Albuterol	Albuterol Medications List	Inhalation
Short-acting, inhaled beta-2 agonists	Levalbuterol	Levalbuterol Medications List	Inhalation

# **Appropriate Testing for Pharyngitis (CWP)**

The CWP measure evaluates the percentage of episodes for patients 3 years and older where the patient was diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode.

## Group A Strep Test Codes:

ш	т и

87070, 87071, 87081, 87430, 87650-87652, 87880

<sup>\*</sup>Codes subject to change

# Follow-Up After Emergency Department Visit for **Substance Use (FUA)**

The FUA measure evaluates the percentage of emergency department (ED) visits among patients age 13 years and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose, for which there was follow-up.

#### Two rates are reported:

- The percentage of ED visits where patients received follow-up within 30 days of the visit (31 total days)
- The percentage of ED visits where patients received follow-up within 7 days of the visit (8 total days)

## 7 & 30 day follow-up codes

Description	Codes
Behavioral Health Assessment	CPT: 99408, 99409 HCPCS: G0396, G0397, G0442, H0001, H0002, H0031, H0049
Online Assessments	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-G2063
Weekly Drug Treatment Service	HCPCS: G2067-G2070, G2072, G2073
Weekly Non Drug Service	HCPCS: G2071, G2074-G2077, G2080
Partial Hospitalization or Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Peer Support Services	HCPCS: G0177, H0024, H0025, H0038-H0040, H0046, H2014, H2023, S9445, T1012, T1016
Substance Use Disorder Services	CPT: 99408, 99409 HCPCS: G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012
Substance Use Services	HCPCS: H0006, H0028
Visit Setting Unspecified include POS code	CPT: 90791, 9072, 90832-90834, 90836- 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99233, 99231- 99233, 99238, 99239, 99251-99255 POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 52, 53, 57, 71, 72
Weekly Drug Treatment Service	HCPCS: G2067-G2070, G2072, G2073
Weekly Non Drug Service	HCPCS: G2071, G2074-G2077, G2080

<sup>\*</sup>Codes subject to change

# Follow-Up After Hospitalization for Mental Illness (FUH)

The FUH measure evaluates percentage of discharges for patients 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health provider.

### Two rates are reported:

- Discharges for which the patient received follow-up within 7 days after discharge
- Discharges for which the patient received follow-up within 30 days after discharge

#### **FUH Measure Codes**

Description	Codes*
Visit Setting Unspecified Value Set with Outpatient POS with Mental Health Provider	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255  POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
BH Outpatient Visit with Mental Health Provider	CPT: 98960-98962, 99078, 99201- 99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381- 99387, 99391-99397, 99401-99404, 99411, 99412, 99492-99494, 99510, 99483 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
Visit Setting Unspecified Value Set with Partial Hospitalization POS with Mental Health Provider	POS: 52
Partial Hospitalization/ Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Visit Setting Unspecified Value Set with Community Mental Health Center POS	POS: 53

## FUH Measure Codes (continued)

Description	Codes*
Electroconvulsive Therapy with	<b>CPT</b> : 90870
Ambulatory Surgical Center POS/	Ambulatory POS: 24
Community Mental Health Center	Comm. POS: 53
POS/ Outpatient POS/ Partial	Partial Hosp. POS: 52
Hospitalization POS	Outpatient POS: 03, 05, 07, 09, 11, 12,
	13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49,
	50, 71, 72
	ICD-10: GZB0ZZZ-GZB4ZZZ
Telehealth Visit POS	CPT: 98966-98968, 99441-99443
	POS: 02
Observation	CPT: 99217-99220
Transitional Care Management	CPT: 99495, 99496
	POS: 53
Psychiatric Collaborative	CPT: 99492-99494
Care Management	HCPCS: G0512

<sup>\*</sup>Codes subject to change

# Follow-Up After Emergency Department Visit for Mental Illness (FUM)

The FUM measure evaluates the percentage of emergency department (ED) visits for patients 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who has a follow-up visit for mental illness.

## Two rates are reported:

- The percentage of ED visits where patients received follow-up within 7 days of the visit (8 total days)
- The percentage of ED visits where patients received follow-up within 30 days of the visit (31 total days)

## 7 & 30 day follow-up codes

Description	Codes
Behavioral Visit	CPT: 98960-98962, 99708, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510  HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-2020, T1015
Electroconvulsive Therapy	CPT: 90870 ICD-10: GZB0ZZZ- ZB4ZZZ
Ambulatory Surgical Center POS	POS: 24
Observation	CPT: 99217-99220
Online Assessment (e-visit or virtual check-in)	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012
Partial Hospitalization or Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Psychiatric Facility-Partial Hospitalization	POS: 52
Telephone Visits	CPT: 98966-98968, 99441-99443
Visit Setting Unspecified with outpatient POS	CPT: 90791, 90792, 90832-90834, 90836- 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255
Outpatient	POS: 02, 03, 05, 07, 09, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 53, 71, 72

<sup>\*</sup>Codes subject to change

# **Initiation and Engagement of Substance Use Disorder Treatment (IET)**

The IET measure evaluates the percentage of new substance use disorder (SUD) episodes for 13 years of age to 64 years of age that result in treatment initiation and engagement:

- Initiation of SUD Treatment: New SUD episodes that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter, partial hospitalization, telehealth visit or medication treatment within 14 days.
- Engagement of SUD Treatment: New SUD episodes that have evidence of treatment engagement within 34 days of initiation.

## Alcohol Use Disorder Treatment Medications

Description	Prescription
Aldehyde dehydrogenase inhibitor	Disulfiram (oral)
Antagonist	Naltrexone (oral and injectible)
Other Acamprosafe (oral; delayed-releatablet)	

### **Opioid Use Disorder Treatment Medications**

Description	Prescription	Medication Lists
Antagonist	Naltrexone (oral)	Naltrexone Oral Medication List
Antagonist	Naltrexone (injectable)	Naltrexone Injection Medication List
Partial Agonist	Buprenorphine (sublingual tablet)	Buprenorphine Oral Medication List
Partial Agonist	Buprenorphine (injection)	Buprenorphine Injection Medication List
Partial Agonist	Buprenorphine (implant)	Buprenorphine Implant Medication List
Partial Agonist	Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)	Buprenorphine Naxolone Medication List

## **IET Medications**

Description	Codes*
BH Outpatient	CPT: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99349, 99350, 99381-99383, 99384-99387, 99391- 99397, 99401- 99404, 99411, 99412, 99483, 99492- 99494, 99510  HCPCS: G0155, G0176, G0177, G0463, G0512, H0002, H0004, H0031, H0034-H0037, H0039, H0040, H0047, H2000, H2010, H2011, H2013-H2020, T1015  Note: Include ICD-10 code for Alcohol abuse and other drug dependence diagnosis
Buprenorphine Implant	HCPCS: G2070, G2072, J0570
Buprenorphine Injection	HCPCS: G2069, Q9991, Q9992
Buprenorphine Naloxone	HCPCS: J0572-J0575
Buprenorphine Oral	HCPCS: H0033, J0571
Buprenorphine Oral Weekly	HCPCS: G2068, F2079
Detoxification	HCPCS: H0008, H0009, H0010-H0014
Methadone Oral	HCPCS: H0020, S0109
Methadone Oral Weekly	HCPCS: G2067, G2078
Naltrexone Injection	HCPCS: G2073, J2315
Online Assessments	CPT: 98969-98972, 99421-99523. 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-G2063
Visit Setting Unspecified that requires Outpatient POS Code	CPT: 90832-90840, 90845, 90847, 90849, 90853, 90875, 908876, 99221- 99223, 99231-99233, 99238, 99239, 99251-99255 POS: 02, 03, 05, 07, 09, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 53, 71, 72
Partial Hospitalization or Intensive Outpatient	HCPCS: H0035, H2001, H2012, S0201, S9480, S9484, S9485
Psychiatric Facility-Partial Hospitalization POS	POS: 52
Substance Use Disorder Services	CPT: 99408, 99409 HCPCS: G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012

Description	Codes*
Monthly Office Based Treatment	HCPCS: G2086, G2087
Weekly Drug Treatment Services	HCPCS: G2067, G2068-G2070, G2072, G2073
Weekly Non Drug Service	HCPCS: G2071, G2074-G2077, G2080

<sup>\*</sup>Codes subject to change

<sup>\*</sup> Codes listed are subject to change, YouthCare recognizes that the circumstances around the services provided may not always directly support/match the codes. It is crucial that the medical record documentation describes the services rendered in order to support the medical necessity and use of these codes.

# **Appendix**

## **Point of Service Code Definitions**

Code	Description
02	Telehealth Provided Other than in Patient's Home
03	School
05	Indian Health Service Free-standing Facility
07	Tribal 638 Free-standing Facility
09	Prison/Correctional Facility
10	Telehealth Provided in Patient's Home
11	Office
12	Home
13	Assisted Living Facility
14	Group Home
15	Mobile Unit
16	Temporary Lodging
17	Walk-in Retail Health Clinic
18	Place of Employment-Worksite
19	Off Campus-Outpatient Hospital
20	Urgent Care Facility
21	Inpatient Hospital
22	On Campus-Outpatient Hospital
23	Emergency Room - Hospital
24	Ambulatory Surgical Center
31	Skilled Nursing Facility
32	Nursing Facility
33	Custodial Care Facility

Code	Description
49	Independent Clinic
50	Federally Qualified Health Center
51	Inpatient Psychiatric Facility
52	Psychiatric Facility-Partial Hospitalization
53	Community Mental Health Center
56	Psychiatric Residential Treatment Center
57	Non-residential Substance Abuse Treatment Facility
58	Non-residential Opioid Treatment Facility
71	Public Health Clinic
72	Rural Health Clinic
81	Independent Laboratory

## **Exclusions**

Exclusions	Applicable Measures	Codes*
Hospice	All Measures	HCPCS: G0182, G9473, G9474, G9475, G9476, G9477, G9478, G9479, Q5003, Q5004, Q5005, Q5006, Q5007, Q5008, Q5010, S9126, T2042, T2043, T2044, T2045, T2046 CPT: 99377, 99378
Palliative Care	BPD, CBP, EED, HBD	HCPCS: G9054, M1017 ICD-10: Z51.5
ESRD Diagnosis	СВР	ICD-10: N18.5-N18.6, Z99.2 ICD-9: 585.5, 585.6, V45.11
Kidney Transplant	СВР	CPT: 50360, 50365, 50380 HCPCS: S2065 ICD-10: OTY00Z0, OTY00Z1, OTY00Z2, OTY10Z0, OTY10Z1, OTY10Z2 ICD-9: 55.61, 55.69
Partial Nephrectomy	СВР	CPT: 50240 ICD-10: OTB00ZZ, OTB03ZZ, OTB04ZZ, OTB07ZZ, OTB08ZZ, OTB10ZZ, OTB13ZZ, OTB14ZZ, OTB17ZZ, OTB18ZZ
Total Nephrectomy	СВР	CPT: 50220, 50225, 50230, 50234, 50236, 50340, 50370, 50543, 50545, 50546, 50548 ICD-10: OTT00ZZ, OTT04ZZ, OTT10ZZ, OTT14ZZ, OTT24ZZ
Fetal Demise	PPC	ICD-10: Z37.1, Z37.4, Z37.7
Polycystic Ovarian syndrome	PPC	ICD-10: E28.2
Gestational Diabetes	PPC, BPD, EED, HBD	ICD-10: O24.410, O24.414, O24.415, O24.419, O24.420, O24.424, O24.425, O24.429, O24.430, O24.434, O24.435, O24.439, O24.911-O24.913, O24.919

Exclusions	Applicable Measures	Codes*
Diabetes Exclusions (including Gestational diabetes, steroid induced diabetes or polycystic ovarian syndrome)		ICD-10: E08.00, E08.01, E08.10, E08.11, E08.21, E08.22, E08.29, E08.311, E08.319, E08.321, E08.3211-E08.3213, E08.3219, E08.329, E08.3291-E08.3293, E08.3299, E08.3311, E08.3313, E08.3319, E08.3311-E08.3313, E08.3319, E08.3399, E08.3391-E08.3393, E08.3399, E08.341, E08.3411-E08.3413, E08.3419, E09.349, E09.3491-E09.3493, E09.3499, E09.3511, E09.3491-E09.3523, E09.3529, E09.3521-E09.3523, E09.3529, E09.3521-E09.3523, E09.3529, E09.3551-E09.3553, E09.3559, E09.3551-E09.3553, E09.3559, E09.3599, E09.3591-E09.3593, E09.3599, E09.36, E09.37X1-E09.37X3, E09.37X9, E09.39-E09.44, E09.49, E09.51-E09.52, E09.59, E09.628, E09.630, E09.638, E09.641, E09.649,
		E09.65, E09.69, E09.8, E09.9, E28.2, O24.410, O24.414-O24.415, O24.419-O24.420, O24.424-O24.425, 429-O24.430, O24.434-O24.435, O24.439, O24.911-O24.913, O24.919, O24.92-O24.93

Exclusions	Applicable	Codes*
<u> </u>	Measures	-coucs
Pregnancy Diagnosis	WCC, CHL* *With a prescription of isotretinoin or an X-Ray within 6 days after pregnancy test	ICD-10: O09.00-O09.03, O09.10-O09.13, O09.211-O09.213, O09.219, O09.291-O09.293, O09.299, O09.30-O09.33, O09.40-O09.43, O09.511-O09.513, O09.519, O09.521-O09.523, O09.529, O09.611-O09.613, O09.621-O09.623, O09.629, O09.70-O09.73, O09.811-O09.813, O09.819, O09.21-O09.823, O09.829, O09.891-O09.893, O09.899, O09.90-O09.93, O09.A0-O09.A3, O10.011-O10.013, O10.019, O10.111-O10.113, O10.119, O10.211-O10.213, O10.219, O10.311-O10.313, O10.319, O10.411-O10.913, O10.919, O11.1, O11.2, O11.3, O11.9, O12.00-O12.03, O12.10-O12.13, O12.20-O12.23, O13.1-O13.3, O13.9, O14.00, O14.02-O14.03, O14.10, O14.12-O14.13, O14.20, O14.22-O14.23, O15.00, O15.02-O15.03, O15.1, O15.9, O16.1-O16.3, O16.9, O20.0, O20.8, O20.9, O21.0-O21.2, O21.8-O21.9, O22.00-O22.03, O22.20-O22.23, O22.30-O22.33, O22.40-O22.43, O22.50-O22.33, O23.10-O23.13, O23.20-O23.03, O23.10-O23.13, O23.20-O23.03, O23.10-O23.13, O23.20-O23.03, O23.10-O23.13, O23.20-O23.03, O23.10-O23.13, O23.599, O23.591-O23.593, O24.011-O24.013, O24.414-O24.415, O24.419, O24.414-O24.813, O24.819, O24.911-O24.913, O24.919, O24.911-O24.913, O24.919, O24.911-O24.913, O24.919, O24.911-O24.913, O24.919, O24.911-O24.913, O24.919, O25.10-O25.13, O26.00-O26.03,

Exclusions	Applicable	Codes*
LACIUSIONS	Measures	Codes
Pregnancy		O26.11-O26.13, O26.20-O26.23,
Diagnosis		026.30-026.33, 026.40-026.43,
(continued)		O26.50-O26.53, O26.611-O26613,
		O26.619, O26.711-O26.713, O26.719,
		O26.811-O26.813, O26.819,
		O26.821-O26.823, O26.829,
		O26.831-O26.833, O26.839,
		O26.841-O26.843, O26.849,
		O26.851-O26.853, O26.859, O26.86,
		O26.872-O26.873, O26.879,
		O26.891-O26.893, O26.899,
		O26.90-O26.93, O28.0-O285,
		O28.8-O28.9, O29.011-O29.013,
		O29.019, O29.021-O29.23, O29.029,
		O29.091-O29.093, O29.099,
		O229.111-O29.113, O29.119,
		O29.121-O29.123, O29.129,
		O29.191-O29.193, O29.199,
		O29.211-O29.213, O29.219,
		O29.291-O29.293, O29.299, O29.3X1-
		O29.3X3, O29.3X9, O29.40-O29.43,
		O29.5X1-O29.5X3, O29.5X9,
		O29.60-O29.63, O29.8X1-O29.8X3,
		O29.8X9, O29.90-O29.93,
		O30.001-O30.003, O30.009,
		O30.011-O30.013, O30.019,
		O30.021-O30.023, O30.029,
		O30.031-O30.033, O30.039,
		O30.041-O30.043, O30.049,
		O30.091-O30-093, O30.099,
		O30.101-O30.103, O30.109,
		O30.111-O30.113, O30.119,
		O30.121-O30.123, O30.129,
		O30.131-O30.133, O30.139,
		O30.191-O30.193, O30.199,
		O30.201-O30.203, O30.209,
		030.211-030.213, 030.219,
		030.221-030.223, 030.229,
		030.231-030.233, 030.239,
		030.291-030.293, 030.299,
		030.801-030.803, 030.809,
		030.811-030.813, 030.819,
		O30.821-O30.823, O30.829,
		O30.831-O30.833, O30.839,

Evalusions	Applicable	Codos*
Exclusions	Applicable Measures	Codes*
Pregnancy		O30.891-O30.893, O30.899,
Diagnosis		O30.90-O30.93, O31.00X0-O30.00X5,
(continued)		O31.00X9, O31.01X0-O31.01X5,
		O31.01X9, O31.02X0-O31.02X5,
		O31.02X9, O31.03X0-O31.03X5,
		O31.03X9, O31.10X0-O31.10X5,
		O31.10X9, O31.11X0-O3111X5,
		O31.11X9, O31.12X0-O31.12X5,
		O31.12X9, O31.13X0-O31.13X5,
		O31.13X9, O31.20X0-O31.20X5,
		O31.20X9, O31.21X0-O31.21X5,
		O31.21X9, O31.22X0-O31.22X5,
		O31.22X9, O31.23X0-O31.23X5,
		O31.23X9, O31.30X0-O31.30X5,
		O31.30X9, O31.31X0-O31.31X5,
		O31.31X9, O31X0-O31.32X5,
		O31.32X9, O33X0-O31.33X5,
		O31.33X9, O31.8X10-O32.8X15,
		O31.8X19, O31.8X20-O31.8X25,
		O31.8X29, O31.8X30-O31.8X35,
		O31.8X39, O31.8X90-O31.8X95,
		O31.8X99, O32.0XX0-O32.0XX5,
		O32.0XX9, O32.1XX0-O32.1XX5,
		O32.1XX9, O32.2XX0-O32.2XX5,
		O32.2XX9, O32.3XX0-O32.3XX5.
		O32.3XX9, O32.4XX0-O32.4XX5,
		O32.4XX9, O32.6XX0-O32.6XX5,
		O32.6XX9, O32.8XX0-O32.8XX5,
		O32.8XX9, O32.9XX0-O32.9XX5,
		O32.9XX9, O33.0-O33.2, O33.3XX0-
		O33.3XX5, O33.3XX9, O33.4XX0-
		O33.4XX5, O33.4XX9, O33.5XX0-
		O33.5XX5, O33.5XX9,
		O33.6XX0-O33.6XX5, O33.6XX9,
		O33.7, O33.7XX0-O33.7XX5, O33.7XX9,
		O33.8, O33.9, 34.00-O34.030,
		034.10-034.13, 034.21,
		034.211-034.212, 034.218-034.219,
		O34.22, O34.29-O34.33,
		034.40-034.43, 034.511-034.513,
		034.519, 034.521-034.523, 034.529,
		O34.531-O34533, O34.539,
		034.591-034.593, 034.599,
		034.60-034.63, 034.70-034.73,
		22

Exclusions	Applicable Measures	Codes*
	Measures	
Pregnancy		034.80-034.83, 034.90-034.93,
Diagnosis		O35.0XX0-O35.0XX5, O35.0XX9,
(continued)		O35.1XX0-O35.1XX5, O35.1XX9,
		O35.2XX0-O35.2XX5, O35.2XX9,
		O35.3XX0-O35.3XX5, O35.3XX9,
		O35.4XX0-O35.4XX5, O35.4XX9,
		O35.5XX0-O35.5XX5, O35.5XX9,
		O35.6XX0-O35.6XX5, O35.6XX9,
		O35.7XX0-O35.7XX5, O35.7XX9,
		O35.8XX0-O35.8XX5, O35.8XX9,
		O35.9XX0-O35.9XX5, O35.9XX9,
		O36.0110-O36.0115, O36.0119,
		O36.0120-O36.0125, O36.0129,
		O36.0130-O36.0135, O36.0139,
		O36.0190-O36.0195, O36.0199,
		O36.0910-O36.0915, O36.0919,
		O36.0920-O36.0925, O36.0929,
		O36.0930-O36.0935, O36.0939,
		O36.0990-O36.0995, O36.0999,
		O36.1110-O36.1115, O36.1119,
		036.1120-036.1125, 036.1129,
		036.1130-036.1135, 036.1139,
		036.1190-036.1195, 036.1199,
		036.1910-036.1915, 036.1919,
		O36.1920-O36.1925, O36.1929,
		O36.1930-O36.1935, O36.1939,
		O36.1990-O36.1995, O36.1999,
		O36.20X0-O36.20X5, O36.20X9,
		O36.21X0-O36.21X5, O36.21X9,
		O36.22X0-O36.22X5, O36.22X9,
		O36.23X0-O36.23X5, O36.23X9,
		O36.4XX0-O36.4XX5, O36.4XX9,
		036.5110-036.5115,
		036.5119-036.5125,
		036.5129-036.5135, 036.5139,
		O36.5190-O36.5195, O36.5199,
		036.5910-036.5915,
		036.5919-036.5925,
		036.5929-036.5935, 036.5939,
		036.5990-036.5995, 036.5999,
		O36.60X0-O36.60X5, O36.60X9,
		O36.61X0-O36.61X5, O36.61X9,
		O36.62X0-O36.621X5, O36.62X9,
		O36.63X0-O36.63X5, O36.63X9,

Exclusions	Applicable	Codes*
Exclusions	Measures	Codes
Pregnancy		O36.70X0-O36.70X5,O36.70X9,
Diagnosis		O36.71X0-O36.71X5, O36.71X9,
(continued)		O36.72X0-O36.72X5, O36.72X9,
		O36.73X0-O36.73X5, O36.73X9,
		O36.80X0-O36.80X5, O36.80X9,
		O36.8120-O36.8125, O36.8129,
		O36.8130-O36.8135, O36.8139,
		O36.8190-O36.8195, O36.8199,
		O36.8210-O36.8215, O36.8219,
		O36.8220-O36.8225, O36.8229,
		O36.8230-O36.8235, O36.8239,
		O36.8290-O36.8295, O36.8299,
		O36.8310-O36.8315, O36.8319,
		O36.8320-O36.8325,
		O36.8329-O36.8335, O36.8339,
		O36.8390-O36.8395, O36.8399,
		O36.8910-O36.8915, O36.8919,
		O36.8920-O36.8925, O36.8929,
		O36.8930-O36.8935, O36.8939,
		O36.8990-O36.8995, O36.8999,
		O36.90X0-O36.90X5, O36.90X9,
		O36.91X0-O36.91X5, O36.91X9,
		O36.92X0-O36.92X5, O36.92X9,
		O36.93X0-O36.93X5, O36.93X9,
		O40.1XX0-O40.1XX5, O40.1XX9,
		O40.2XX0-O40.2XX5, O40.2XX9,
		O4.31XX0-O40.3XX5, O40.3XX9,
		O40.9XX0-O40.9XX5, O40.9XX9,
		O41.00X0-O41.00X5, O41.00X9,
		O41.01X0-O41.01X5, O41.01X9,
		O41.02X0-O41.02X5, O41.02X9,
		O41.03X0-O41.03X5, O41.03X9,
		O41.1010-O41.1015,
		O41.1019-O41.1025,
		O41.1029-O41.1035, O41.1039,
		O41.1090-O41.1095, O41.1099,
		O41.1210-O41.1215,
		O41.1219-O41.1225,
		O41.1229-O41.1235, O41.1239,
		O41.1290-O41.1295, O41.1299,
		O41.1410-O41.1415,
		O41.1419-O41.1425,
		O41.1429-O41.1435, O41.1439,
		O41.1490-O41.1495, O41.1499,

E. A. Maria	A P L.L.	Color
Exclusions	Applicable Measures	Codes*
Pregnancy	1 reasures	O41.8X10-O41.8X15, O41.8X19-
Diagnosis		O41.8X25, O41.8X29-O41.8X35,
(continued)		O41.8X39, O41.8X91-O41.8X95,
(Continued)		O41.8X99, O41.90X0-O41.90X5,
		O41.90X9, O41.91X0-O41.91X5,
		O41.91X9, O41.92X0-O41.92X5,
		O41.92X9, O41.93X0-O41.93X5,
		O41.93X9, O42.00, O42.011-O42.013,
		O42.019, O42.02, O42.10,
		O42.111-O42.113, O42.119, O42.12,
		O42.011-O42.013, O42.113, O42.12,
		O42.911-O42.913, O42.919, O42.92,
		O43.011-O43.013, O43.019,
		O43.021-O43.023, O43.029,
		O43.101-O43.103, O43.109,
		O43.111-O43.113, O43.119, O43.121-O43.123. O43.129,
		•
		O43.191-O43.193, O43.199,
		043.211-043.213, 043.219,
		O43.221-O43.223, O43.229,
		O43.231-O43.233, O43.239,
		043.811-043.813, 043.819,
		O43.891-O43.893, O43.899,
		043.90-043.93, 044.00-044.03,
		044.10-044.13, 044.20-044.23,
		044.30-044.33, 044.40-044.43,
		O44.50-O44.53, O45.001-O45.003,
		O45.009, O45.011-O45.013, O45.019,
		O45.021-O45.023, O45.029,
		O45.091-O45.093, O45.099, O45.8X1-
		O45.8X3, O45.8X9, O45.90-O45.93,
		O46.001-O46.003, O46.009,
		046.011-046.013, 046.019,
		O46.021-O46.023, O46.029,
		046.091-046.093, 046.099,
		O46.8X1-O46.8X3, O46.8X9,
		046.90-046.93, 047.00-047.03,
		047.1, 047.9, 048.0-048.1, 060.00,
		060.02-060.03, 071.00-
		071.02-071.03, 071.1-071.7,
		071.81-071.82, 071.89, 071.9,
		088.011-088.013, 088.019,
		O88.111-O88.113, O88.119,
		O88.211-O88.213, O88.219,

Exclusions	Applicable Measures	Codes*
Pregnancy		088.311-088.313, 088.319,
Diagnosis		088.811-088.813, 088.819,
(continued)		O91.011-O91.013, O91.019, O91.13,
		O91.211-O91.213, O91.219, O91.23,
		O92.011-O92.013, O92.019, O92.03,
		O92.111-O92.113, O92.119,
		092.3-092.6, 092.70, 092.79,
		O98.011-O98.013, O98.019,
		O98.111-O98.113, O98.119,
		O98.211-O98.213, O98.219,
		O98.311-O98.313, O98.319,
		O98.411-O98.413, O98.419,
		O98.511-O98.513, O98.519,
		O98.611-O98.613, O98.619,
		O98.711-O98.713, O98.719,
		O98.811-O98.813, O98.819,
		O98.911-O98.913, O98.919,
		O99.011-O99.013, O99.019,
		O99.111-O99.113, O99.119,
		O99.210-O99.213, O99.280-O99.283,
		O99.310-O99.313, O99.320-O99.323,
		099.330-099.333, 099.340-099.343,
		O99.350-O99.353, O99.411-O99.413,
		O99.419, O99.511-O99.513, O99.519,
		O99.611-O99.613, O99.619,
		O99.711-O99.713, O99.719, O99.810,
		O99.820, O99.830, O99.840-O99.843,
		O99.891, O9A.111-O9A.113, O9A.119,
		O9A.211-O9A.213, O9A.219, O9A.311-
		O9A.313, O9A.319, O9A.411-O9A.413,
		O9A.419, O9A.511-O9A.513, O9A.519,
		Z03.71-Z03.75, Z03.79, Z32.01,
		Z34.00-Z34.03, Z34.80-Z34.83,
		Z34.90-Z34.93, Z36, Z36.0-Z36.5,
		Z36.81-Z36.89, Z36.8A, Z36.9

Exclusions	Applicable Measures	Codes*
Exclusions  Diagnostic Radiology	Applicable Measures CHL* *With a dx of pregnancy	Codes*  CPT: 70010, 70015, 70030, 70100, 70110, 70120, 70130, 70134, 70140, 70150, 70160, 70170, 70190, 70200, 70210, 70220, 70240, 70260, 70300, 70310, 70320, 70328, 70330, 70332, 70336, 70350, 70355, 70360, 70370, 70371, 70380, 70390, 70450, 70460, 70470, 70492, 70496, 70498, 70540, 70542-70549, 70551-70559, 71045-7148, 71100-71101, 71110-71111, 71120, 71130, 71250, 71260, 71270-71271, 71275, 71550-71552, 71555, 72020, 72040, 72050, 72052, 72070, 72072, 72074, 72880-72084, 72100, 72110, 72114, 72120, 7215-72133, 72141-72142, 72147-72149, 72156-72159, 72170, 72190-72198, 72200, 72002, 72202, 72220, 72240, 72255, 72265, 72270, 72275, 72285, 72295, 73000, 73010, 73020, 73030, 73040, 07050, 73060, 73070, 73080, 73088, 73090, 73092, 73100, 73110, 73115, 73120, 73130, 73140, 73200-73202, 73206, 73218-73223, 73225, 73561-73552, 73560, 73562, 73564-73565, 73580, 73590, 73592, 73600, 73610, 73615, 73620, 73630, 73660, 73610, 73615, 73620, 7378-73723, 73725, 73018-73723, 73725, 74018-74019, 74021-74022, 74150, 74160, 74170, 74174-74178, 74181-74183, 74180, 74280, 74283, 74290, 74355, 74360, 74330, 74340, 74355, 74360, 74330, 74340, 74355, 74360, 74363, 74400, 74415, 74420, 74425, 74430, 74445, 74440, 74445, 74440, 74475, 74450, 74455, 74450, 74460, 74475, 74450, 74445, 74450, 74455, 74450, 74460, 74475, 74450, 74460, 74475, 74440, 74475, 74476, 7
		74440, 74445, 74450, 74455, 74470, 74485, 74710, 74712-74713, 74740, 74742, 74775, 75557, 75559, 75561, 75563, 75565, 75571-75574, 75600,
		756005, 75625, 75630, 75635, 75705,

Exclusions	Applicable Measures	Codes*
Diagnostic Radiology (continued)		75710, 75716, 75726, 75731, 75733, 75736, 75741, 75743, 75746. 75756, 75774, 75801, 75803, 75805, 75807, 75809-75810, 75820, 75822, 75825, 75827, 75831, 75833, 75840, 75842, 75860, 75870,75872, 75880, 75885, 75887, 75889, 75891, 75893-75894, 75898, 75901-75902, 75956-75959, 75970, 75984, 75989, 76000, 76010, 76080, 760998, 76100-76102, 76120, 76125, 76140, 76145, 76376-76377, 76380, 76390-76391, 76496-76499
Severe Combined Immunodeficiency	CIS	ICD-10: D81.0-D81.2, D81.9
Disorders of the Immune System	CIS	ICD-10: D80.0-D80.9, D81.0-D81.4, D81.6, D81.6-D81.7, D81.89, D81.9, D82.0-D82.4, D82.8-D82.9, D83.0-D83.2, D83.8-D83.9, D84.0-D84.1, D84.8, D84.81, D84.821-D84.822, D84.89, D84.9, D89.3, D89.810-D89.813, D89.52, D89.831-D89.835, D89.835, D89.839, D89.89, D89.9
HIV	CIS	ICD-10: B20, Z21
HIV Type 2	CIS	ICD-10: B97.35

Exclusions	Applicable Measures	Codes*
Malignant Neoplasm of Lymphatic Tissue	CIS	ICD-10: C81.00-C81.49, C81.71-C81.79, C81.90-C81.99, C82.00-C82.39, C83.51-C83.59, C83.70-C83.99, C84.00-C84.19, C84.40-C84.49, C84.60-C84.29, C85.10-C85.29, C85.80-C85.99, C86.0-C86.6, C88.2-C88.9, C90.00-C90.02, C91.10-C90.12, C90.20-C90.22, C90.30-C90.32, C91.00-C91.02, C91.10-C91.12, C91.30-C91.32, C94.40-C91.42, C91.50-C91.52, C91.60-C91.62, C91.90-C91.22, C92.00-C92.02, C92.10-C92.12, C92.00-C92.02, C92.10-C92.12, C92.00-C92.02, C92.30-C92.32, C92.40-C92.42, C92.50-C92.52, C92.60-C92.62, C92.90-C92.92, C92.A0-C93.32, C93.90-C93.92, C93.30-C93.32, C93.90-C93.92, C93.30-C93.32, C94.00-C94.02, C94.20-C94.22, C94.30-C94.32, C94.80-C94.82, C95.00-C95.02, C95.10-C95.12, C95.90-C95.92, C96.0, C96.2, C96.20-C96.22, C96.29, C96.4, C96.9, C96.A, C96.Z
Intussusception	CIS	ICD-10: K56.1
Narcolepsy	ADD	ICD-10: G47.411, G47.419, G47.421, G47.429