

# Clinical Policy: Measurement of Serum 1,25-dihydroxyvitamin D

Reference Number: CP.MP.152

Last Review Date: 09/2022

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

## Description

Vitamin D is metabolized in the liver to 25-hydroxyvitamin D [25(OH)D], (also known as calcidiol), and then in the kidney to 1,25-dihydroxyvitamin D [1,25(OH)2D], also known as calcitriol. 25(OH)D is the major circulating form of vitamin D while 1,25(OH)2D is the active form of vitamin D. In individuals at risk for vitamin D deficiency, the best method for determining a person's vitamin D status is to measure a 25(OH)D concentration. Measurement of 1,25(OH)2D is not useful for monitoring the vitamin D status, as it does not reflect vitamin D reserves.<sup>1</sup> This policy addresses when measurement of 1,25(OH)2D is appropriate and medically necessary.

## Policy/Criteria

- I. It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that measurement of serum 1,25(OH)2D (CPT 82652) is **medically necessary** for monitoring certain conditions, such as acquired and inherited disorders of vitamin D and phosphate metabolism, including any of the following indications:
  - A. Chronic kidney disease;
  - B. Hereditary phosphate-losing disorders;
  - C. Oncogenic osteomalacia;
  - D. Pseudovitamin D-deficiency rickets;
  - E. Vitamin D-resistant rickets;
  - F. Chronic granuloma-forming disorders (e.g., sarcoidosis and some lymphomas).
  
- II. It is the policy of health plans affiliated with Centene Corporation that measurement of serum 1,25(OH)2D for routine screening of average risk, asymptomatic individuals is **not medically necessary**.

## Background

Vitamin D or calciferol, is a fat-soluble vitamin that plays an important role in calcium homeostasis and bone health. Vitamin D comes in two forms, D<sub>2</sub> and D<sub>3</sub>. It is unique among hormones because the major source of vitamin D is exposure to natural sunlight. Very few foods naturally contain, or are fortified with, vitamin D, thus, the major cause of vitamin D deficiency is inadequate exposure to sunlight.

The optimal serum 25(OH)D concentration for skeletal health is controversial, however, experts agree that levels lower than 20 ng/mL are suboptimal for skeletal health.<sup>5</sup> Vitamin D deficiency is defined by the Endocrine Society as a 25(OH)D below 20 ng/ml (50 nmol/liter). Vitamin D deficiency results in abnormalities in calcium, phosphorus, and bone metabolism. It causes a decrease in the efficiency of intestinal calcium and phosphorus absorption of dietary calcium and phosphorus, resulting in an increase in parathyroid hormone (PTH) levels. Secondary

## CLINICAL POLICY

### Measurement of Serum 1,25-dihydroxyvitamin D

hyperparathyroidism maintains serum calcium in the normal range at the expense of mobilizing calcium from the skeleton and increasing phosphorus wasting in the kidneys.

Screening for Vitamin D deficiency is recommended for individuals at risk, such as those with osteomalacia, osteoporosis, chronic kidney disease, hepatic failure, malabsorption syndromes, hyperparathyroidism, African-American and Hispanic children and adults, pregnant or lactating women, older adults with history of falls or non-traumatic fractures, obese children or adults (BMI greater than 30 kg/m<sup>2</sup>), granuloma-forming disorders, and some lymphomas.<sup>1</sup>

Circulating 25(OH)D is the best indicator to monitor for vitamin D status as it is the main circulating form of vitamin D, and has a half-life of two to three weeks. In contrast, 1,25(OH)2D, has a much shorter half-life of about four hours, circulates in much lower concentrations than 25(OH)D, and is susceptible to fluctuations induced by PTH in response to subtle changes in calcium levels. Serum 1,25(OH)2D is frequently either normal or even elevated in those with vitamin D deficiency, due to secondary hyperparathyroidism.<sup>1</sup>

#### *The Endocrine Society*

The Endocrine Society recommends using the serum circulating 25-hydroxyvitamin D [25(OH)D] level, measured by a reliable assay, to evaluate vitamin D status in patients who are at risk for vitamin D deficiency and in whom a prompt response to optimization of vitamin D status could be expected. They note further, 1,25(OH)2D measurement does not reflect vitamin D status as levels are tightly regulated by serum levels of PTH, calcium, and phosphate. Serum 1,25(OH)2D does not reflect vitamin D reserves, and measurement of 1,25(OH)2D is not useful for monitoring the vitamin D status of patients. Serum 1,25(OH)2D is frequently either normal or even elevated in those with vitamin D deficiency, due to secondary hyperparathyroidism. Measurement of 1,25(OH)2D is useful in acquired and inherited disorders in the metabolism of 25(OH)D and phosphate, including chronic kidney disease, hereditary phosphate-losing disorders, oncogenic osteomalacia, pseudovitamin D-deficiency rickets, vitamin D-resistant rickets, as well as chronic granuloma-forming disorders such as sarcoidosis and some lymphomas.

#### *United States Preventive Services Task Force (USPSTF)*

The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening for vitamin D deficiency in asymptomatic adults.

#### *American Congress of Obstetricians and Gynecologists*

At this time, there is insufficient evidence to support a recommendation for screening all pregnant women for vitamin D deficiency. For pregnant women thought to be at increased risk of vitamin D deficiency, maternal serum 25-hydroxyvitamin D levels can be considered and should be interpreted in the context of the individual clinical circumstance.<sup>3</sup>

### **Coding Implications**

This clinical policy references Current Procedural Terminology (CPT<sup>®</sup>). CPT<sup>®</sup> is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2019, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are

**CLINICAL POLICY**

**Measurement of Serum 1,25-dihydroxyvitamin D**

included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
82652	Vitamin D; 1, 25 dihydroxy, includes fraction(s), if performed

HCPCS Codes	Description
N/A	

**ICD-10-CM Diagnosis Codes that Support Coverage Criteria**

ICD-10-CM Code	Description
A15.0-A15.9	Respiratory tuberculosis
C81.00-C81.99	Hodgkin lymphoma
C82.00-C82.99	Follicular lymphoma
C83.00-C83.99	Non-follicular lymphoma
C84.00-C84.99	Mature T/NK-cell lymphomas
C88.0-C88.9	Malignant immunoproliferative diseases and certain other B-cell lymphomas
D86.0-D86.9	Sarcoidosis
E20.0	Idiopathic hypoparathyroidism
E20.8	Other hypoparathyroidism
E21.0-E21.5	Hyperparathyroidism and other disorders of parathyroid gland
E55.0	Rickets, active
E83.30-E83.39	Disorder of phosphorus metabolism and phosphatases
E83.50-E83.59	Disorders of calcium metabolism
N18.1-N18.9	Chronic kidney disease (CKD)
N25.0	Renal osteodystrophy
P37.0	Congenital tuberculosis

Reviews, Revisions, and Approvals	Date	Approval Date
Policy developed	11/17	12/17
Removed CPT code 82306 as the policy does not apply to this test.	09/18	
References reviewed and updated	10/18	11/18
References reviewed and updated. Code E20.00 corrected to E20.0.	11/19	11/19

**References**

- Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, Treatment, and Prevention of Vitamin D Deficiency: an Endocrine Society Clinical Practice Guideline. The Journal of Clinical Endocrinology & Metabolism, Volume 96, Issue 7, 1 July 2011, Pages 1911–1930.

**CLINICAL POLICY****Measurement of Serum 1,25-dihydroxyvitamin D**

Accessed Oct 24 2018. Available at: <https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2011-0385>

2. United States Preventive Services Task Force. Vitamin D Deficiency: Screening November 2014
3. American College of Obstetricians and Gynecologists. Vitamin D: Screening and Supplementation During Pregnancy. Committee Opinion Number 495. July 2011. Reaffirmed 2019.
4. Pazirandeh S, Burns DL. Overview of vitamin D. In: UpToDate, Waltham, MA. Motil KJ, Drezner MK (Eds). Accessed 11/6/19
5. Dawson-Hughes B. Vitamin D deficiency in adults: Definition, clinical manifestations, and treatment. In: UpToDate, Waltham, MA. Drezner MK, Rosen CJ (Eds). Accessed 11/6/19.
6. Misra M. Vitamin D insufficiency and deficiency in children and adolescents. In: UpToDate, Waltham, MA. Motil KJ, Drezner MK (Eds). Accessed 11/6/19.
7. Drezner MK. Causes of vitamin D deficiency and resistance. In: UpToDate, Waltham, MA. Rosen CJ SA (Ed). Accessed 11/6/19.
8. Tebben PJ, Singh RJ, Kumar R. Vitamin D-Mediated Hypercalcemia: Mechanisms, Diagnosis, and Treatment. *Endocr Rev.* 2016 Oct;37(5):521-547. Epub 2016 Sep 2.
9. Florenzano P, Gafni RI, Collins MT. Tumor-induced osteomalacia. *Bone Rep.* 2017 Sep 20;7:90-97. doi: 10.1016/j.bonr.2017.09.002. eCollection 2017 Dec.
10. Ruppe MD. X-Linked Hypophosphatemia. *GeneReviews*<sup>®</sup> [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2017. 2012 Feb 9 [updated 2017 Apr 13].

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a

## CLINICAL POLICY

### Measurement of Serum 1,25-dihydroxyvitamin D

discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

**Note: For Medicare members**, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.