CLINICAL POLICY

Ferric Maltol



Clinical Policy: Ferric Maltol (Accrufer)

Reference Number: IL.PMN.213

Effective Date: 09.03.19 Last Review Date: 04.08.22 Line of Business: Medicaid

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Ferric maltol (Accrufer[™]) is an iron replacement product.

FDA Approved Indication(s)

Accrufer is indicated for the treatment of iron deficiency in adults.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Accrufer is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- **A. Iron Deficiency** (must meet all):
 - 1. Diagnosis of iron deficiency;
 - 2. Age \geq 18 years;
 - 3. Failure of two oral iron products (*must be different salts*), unless clinically significant adverse effects are experienced or contraindicated;
 - 4. Dose does not exceed 60 mg (2 capsules) per day.

Approval duration: 12 months

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized CP.PMN.53 for Medicaid.

II. Continued Therapy

- **A. Iron Deficiency** (must meet all):
 - 1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - 2. Member is responding positively to therapy;
 - 3. If request is for a dose increase, new dose does not exceed 60 mg (2 capsules) per day.

Approval duration: 12 months



B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 12 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
ferrous fumarate (Ferrimin 150,	PO; dose and frequency varies	Varies
Ferretts, Ferrocite, Hemocyte)	DO: dose and frequency varies	Varias
ferrous gluconate (Fergon, Ferrotabs) ferrous sulfate (Feosol, Ferro-Bob,	PO: dose and frequency varies	Varies Varies
FerrouSul)	PO; dose and frequency varies	varies
polysaccharide-iron complex (EZFE 200, Ferrex 150, Ferric-X 150, iFerex	PO; dose and frequency varies	Varies
150, Myferon 150, NovaFerrum 50,		
Nu-iron 150, PIC 200, Poly-Iron 150)		

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity to the active substance or any excipient; hemochromatosis and other iron overload syndromes; patients receiving repeated blood transfusions
- Boxed warning(s): none reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Iron	30 mg PO BID, taken 1 hour before or 2 hours after a	60 mg/day
deficiency	meal	

CLINICAL POLICY Ferric Maltol



Indication	Dosing Regimen	Maximum Dose
	Treatment duration will depend on the severity of iron deficiency but generally at least 12 weeks of treatment is required. The treatment should be continued as long as necessary until ferritin levels are within the normal range	

VI. Product Availability

Capsule: 30 mg

VII. References

 Accrufer Prescribing Information. London: Shield Therapeutics; March 2022. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/212320Orig1s000lbl.pdf. Accessed April 8, 2022.

Reviews, Revisions, and Approvals	Date	P&T Approval
		Date
Policy created	09.03.19	11.19
Removed references to Centene preferred product verbiage for	12.11.19	
migration to HFS PDL.		
Removed references to HIM/Commercial, Medicaid policy only.		
4Q 2020 annual review: no significant changes	12.15.20	
2Q 2021 annual review:	6.30.21	
added table Appendix B: Therapeutic Alternatives. Reviewed and		
updated references		
2Q 2022 annual review: Reviewed and updated references	4.8.22	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage

CLINICAL POLICY Ferric Maltol



decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2019 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.