

## Clinical Policy: Tofacitinib (Xeljanz, Xeljanz XR)

Reference Number: IL.PHAR.267

Effective Date: 12/13/2021

Last Review Date: 12/13/2021

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Tofacitinib (Xeljanz<sup>®</sup>, Xeljanz<sup>®</sup> XR) is a Janus kinase (JAK) inhibitor.

### FDA Approved Indication(s)

Xeljanz and Xeljanz XR are indicated for the treatment of:

- Adult patients with moderately to severely active rheumatoid arthritis (RA) who have had an inadequate response or intolerance to methotrexate (MTX).
- Adult patients with active psoriatic arthritis (PsA) who have had an inadequate response or intolerance to MTX or other disease-modifying antirheumatic drugs (DMARDs).
- Adult patients with moderately to severely active ulcerative colitis (UC) who have had an inadequate response or who are intolerant to tumor necrosis factor (TNF) blockers.

Adult patients with active ankylosing spondylitis who have had an inadequate response or intolerance to one or more TNF blockers.

Xeljanz is additionally indicated for active polyarticular course juvenile idiopathic arthritis (pcJIA) in patients 2 years of age and older who have had an inadequate response or intolerance to one or more TNF blockers.

Limitation(s) of use: Use of Xeljanz/Xeljanz XR in combination with biologic DMARDs or potent immunosuppressants such as azathioprine and cyclosporine is not recommended.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Xeljanz and Xeljanz XR are **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Polyarticular Course Juvenile Idiopathic Arthritis (must meet all):

1. Diagnosis of pcJIA as evidenced by  $\geq 5$  joints with active arthritis;
2. Request is for Xeljanz immediate-release tablets or oral solution;
3. Prescribed by or in consultation with a rheumatologist;
4. Age  $\geq 2$  years;
5. Documented baseline 10-point clinical juvenile arthritis disease activity score (cJADAS-10) (*see Appendix I*);
6. Member meets one of the following (a, b, c, or d):

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

- a. Failure of a  $\geq 3$  consecutive month trial of MTX at up to maximally indicated doses;
  - b. Member has intolerance or contraindication to MTX (*see Appendix D*), and failure of a  $\geq 3$  consecutive month trial of leflunomide or sulfasalazine at up to maximally indicated doses, unless clinically significant adverse effects are experienced or both are contraindicated;
  - c. For sacroiliitis/axial spine involvement (i.e., spine, hip), failure of a  $\geq 4$  week trial of an NSAID at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
  - d. Documented presence of high disease activity as evidenced by cJADAS-10  $> 8.5$  (*see Appendix I*);
7. Member has not responded or is intolerant to one or more TNF blockers, unless contraindicated;  
*\*Prior authorization may be required for TNF blockers*
8. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
9. Dose does not exceed 10 mg (2 tablets or 10 mL) per day.

**Approval duration: 6 months**

#### **B. Psoriatic Arthritis (must meet all):**

1. Diagnosis of PsA;
2. Prescribed by or in consultation with a dermatologist or rheumatologist;
3. Age  $\geq 18$  years;
4. Member has not responded or is intolerant to one or more TNF blockers, unless contraindicated;  
*\*Prior authorization may be required for TNF blockers*
5. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
6. Dose does not exceed one of the following (a or b):
  - a. Xeljanz: 10 mg (2 tablets) per day;
  - b. Xeljanz XR: 11 mg (1 tablet) per day.

**Approval duration: 6 months**

#### **C. Rheumatoid Arthritis (must meet all):**

1. Diagnosis of RA per American College of Rheumatology (ACR) criteria (*see Appendix F*);
2. Prescribed by or in consultation with a rheumatologist;
3. Age  $\geq 18$  years;
4. Member meets one of the following (a or b):
  - a. Failure of a  $\geq 3$  consecutive month trial of MTX at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
  - b. If intolerance or contraindication to MTX (*see Appendix D*), failure of a  $\geq 3$  consecutive month trial of at least ONE conventional DMARD (e.g., sulfasalazine, leflunomide, hydroxychloroquine) at up to maximally indicated

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

doses, unless clinically significant adverse effects are experienced or all are contraindicated;

5. Documentation of one of the following baseline assessment scores (a or b):
  - a. Clinical disease activity index (CDAI) score (*see Appendix G*);
  - b. Routine assessment of patient index data 3 (RAPID3) score (*see Appendix H*);
6. Member has not responded or is intolerant to one or more TNF blockers, unless contraindicated;
7. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
8. Dose does not exceed one of the following (a or b):
  - a. Xeljanz: 10 mg (2 tablets) per day;
  - b. Xeljanz XR: 11 mg (1 tablet) per day.

**Approval duration: 6 months**

#### **D. Ulcerative Colitis (must meet all):**

1. Diagnosis of moderately to severely UC;
2. Prescribed by or in consultation with a gastroenterologist;
3. Age  $\geq$  18 years;
4. Member has not responded or is intolerant to one or more TNF blockers, unless contraindicated;
5. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
6. Dose does not exceed one of the following (a or b):
  - a. Xeljanz: 20 mg (2 tablets) per day;
  - b. Xeljanz XR: 22 mg (1 tablet) per day.

**Approval duration: 6 months**

#### **E. Ankylosing Spondylitis (must meet all):**

1. Diagnosis of AS;
2. Prescribed by or in consultation with a rheumatologist;
3. Age  $\geq$  18 years;
4. Failure of at least TWO non-steroidal anti-inflammatory drugs (NSAIDs) at up to maximally indicated doses, each used for  $\geq$  4 weeks unless contraindicated or clinically significant adverse events are experienced;
5. Member has not responded or is intolerant to one or more TNF blockers, unless contraindicated;  
*\*Prior authorization may be required for TNF blockers*
6. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
7. Dose does not exceed one of the following (a or b):
  - a. Xeljanz (both i and ii):
    - i. 10 mg per day;
    - ii. 2 tablets per day;

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

- b. Xeljanz XR (both i and ii):
  - i. 11 mg per day;
  - ii. 1 tablet per day.

#### **Approval duration: 6 months**

#### **F. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.PMN.53 for Medicaid.

## **II. Continued Therapy**

### **A. All Indications in Section I (must meet all):**

1. Member meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member meets one of the following (a, b, or c):
  - a. For RA: Member is responding positively to therapy as evidenced by one of the following (i or ii):
    - i. A decrease in CDAI (*see Appendix G*) or RAPID3 (*see Appendix H*) score from baseline;
    - ii. Medical justification stating inability to conduct CDAI re-assessment, and submission of RAPID3 score associated with disease severity that is similar to initial CDAI assessment or improved;
  - b. For psJIA: Member is responding positively to therapy as evidenced by a decrease in cJADAS-10 from baseline (*see Appendix I*);
  - c. For all other indications: Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed one of the following (a or b):
  - a. Xeljanz (i, ii, or iii):
    - i. RA or PsA: 10 mg (2 tablets) per day;
    - ii. UC: 20 mg (2 tablets) per day;

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

- iii. pcJIA: 10 mg (2 tablets or 10 mL) per day;
- b. Xeljanz XR (i or ii):
  - i. RA or PsA: 11 mg (1 tablet) per day;
  - ii. UC: 22 mg (1 tablet) per day.

#### **Approval duration: 12 months**

#### **B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.PMN.53 for Medicaid.

#### **III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 for Medicaid or evidence of coverage documents;
- B. Combination use with biological disease-modifying antirheumatic drugs (bDMARDs) or potent immunosuppressants, including but not limited to any tumor necrosis factor (TNF) antagonists [e.g., Cimzia<sup>®</sup>, Enbrel<sup>®</sup>, Humira<sup>®</sup> and its biosimilars, Simponi<sup>®</sup>, Avsola<sup>™</sup>, Inflectra<sup>™</sup>, Remicade<sup>®</sup>, Renflexis<sup>™</sup>], interleukin agents [e.g., Arcalyst<sup>®</sup> (IL-1 blocker), Ilaris<sup>®</sup> (IL-1 blocker), Kineret<sup>®</sup> (IL-1RA), Actemra<sup>®</sup> (IL-6RA), Kevzara<sup>®</sup> (IL-6RA), Stelara<sup>®</sup> (IL-12/23 inhibitor), Cosentyx<sup>®</sup> (IL-17A inhibitor), Taltz<sup>®</sup> (IL-17A inhibitor), Siliq<sup>™</sup> (IL-17RA), Ilumya<sup>™</sup> (IL-23 inhibitor), Skyrizi<sup>™</sup> (IL-23 inhibitor), Tremfya<sup>®</sup> (IL-23 inhibitor)], Janus kinase inhibitors (JAKi) [e.g., Xeljanz<sup>®</sup>/Xeljanz<sup>®</sup> XR, Cibinco<sup>™</sup>, Olumiant<sup>™</sup>, Rinvoq<sup>™</sup>], anti-CD20 monoclonal antibodies [Rituxan<sup>®</sup>, Riabni<sup>™</sup>, Ruxience<sup>™</sup>, Truxima<sup>®</sup>, Rituxan Hycela<sup>®</sup>], selective co-stimulation modulators [Orencia<sup>®</sup>], and integrin receptor antagonists [Entyvio<sup>®</sup>] because of the additive immunosuppression, increased risk of neutropenia, as well as increased risk of serious infections;
- C. Alopecia areata (ICD10: L63), also referred to as patchy hairloss.

#### **IV. Appendices/General Information**

##### *Appendix A: Abbreviation/Acronym Key*

CDAI: clinical disease activity index  
cJADAS: clinical juvenile arthritis  
disease activity score

DMARDs: disease-modifying  
antirheumatic drugs  
FDA: Food and Drug Administration

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

JAK: Janus kinase  
 MTX: methotrexate  
 pcJIA: polyarticular course juvenile  
 idiopathic arthritis;  
 RA: rheumatoid arthritis

RAPID3: routine assessme of patient index  
 data 3  
 PsA: psoriatic arthritis  
 UC: ulcerative colitis

#### Appendix B: Therapeutic Alternatives

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
azathioprine (Azasan <sup>®</sup> , Imuran <sup>®</sup> )	<b>RA</b> 1 mg/kg/day PO QD or divided BID	2.5 mg/kg/day
Cuprimine <sup>®</sup> (d-penicillamine)	<b>RA*</b> <u>Initial dose:</u> 125 or 250 mg PO QD <u>Maintenance dose:</u> 500 – 750 mg/day PO QD	1,500 mg/day
corticosteroids	<b>UC*</b> <i>Adult:</i> Prednisone 40 mg – 60 mg PO QD, then taper dose by 5 to 10 mg/week  Budesonide (Uceris <sup>®</sup> ) 9 mg PO QAM for up to 8 weeks	Various
cyclosporine (Sandimmune <sup>®</sup> , Neoral <sup>®</sup> )	<b>RA</b> 2.5 – 4 mg/kg/day PO divided BID	4 mg/kg/day
hydroxychloroquine (Plaquenil <sup>®</sup> )	<b>RA*</b> <u>Initial dose:</u> 400 – 600 mg/day PO QD <u>Maintenance dose:</u> 200 – 400 mg/day PO QD	600 mg/day
leflunomide (Arava <sup>®</sup> )	<u>Initial dose (for low risk hepatotoxicity or myelosuppression):</u> 100 mg PO QD for 3 days <u>Maintenance dose:</u> 20 mg PO QD  <b>PJIA*</b> Weight < 20 kg: 10 mg every other day Weight 20 - 40 kg: 10 mg/day Weight > 40 kg: 20 mg/day	20 mg/day

**CLINICAL POLICY**  
**Tofacitinib/Tofacitinib XR**

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
methotrexate (Trexall <sup>®</sup> , Otrexup <sup>™</sup> , Rasuvo <sup>®</sup> , RediTrex <sup>®</sup> , Xatmep <sup>™</sup> , Rheumatrex <sup>®</sup> )	<b>RA</b> 7.5 mg/week PO, SC, or IM or 2.5 mg PO Q12 hr for 3 doses/week  <b>PJIA*</b> 10 – 20 mg/m <sup>2</sup> /week PO, SC, or IM	30 mg/week
Ridaura <sup>®</sup> (auranofin)	<b>RA</b> 6 mg PO QD or 3 mg PO BID	9 mg/day (3 mg TID)
sulfasalazine (Azulfidine <sup>®</sup> )	<b>RA</b> <u>Initial dose:</u> 500 mg to 1,000 mg PO QD for the first week. Increase the daily dose by 500 mg each week up to a maintenance dose of 2 g/day. <u>Maintenance dose:</u> 2 g/day PO in divided doses  <b>PJIA*</b> 30-50 mg/kg/day PO divided BID	RA: 3 g/day  PJIA: 2 g/day
NSAIDs (e.g., indomethacin, ibuprofen, naproxen, celecoxib)	<b>PJIA*</b> Varies	Varies

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*\*Off-label*

**Appendix C: Contraindications/Boxed Warnings**

- Contraindication(s): none reported
- Boxed warning(s):
  - Serious infections: There is an increased risk of serious infections leading to hospitalization or death.
  - Mortality: Rheumatoid arthritis patients 50 years and older with at least one cardiovascular risk factor treated with Xeljanz 10 twice daily had a higher rate of all-cause mortality, including sudden CF death, compared to those treated with Xeljanz 5 mg given twice daily or TNF blockers in a large, ongoing, post marketing study.
  - Malignancies: Lymphoma and other malignancies, as well as Epstein Barr Virus-associated post-transplant lymphoproliferative disorder has been observed.
  - Thrombosis: Thrombosis, including pulmonary embolism, deep venous thrombosis, and arterial thrombosis have occurred in patients treated with Xeljanz and other Janus kinase inhibitors used to treat inflammatory conditions.
    - A large, ongoing postmarketing safety study observed an increase in incidence of thrombosis events in rheumatoid arthritis patients who were 50 years of age and

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

older with at least one CV risk factor treated with Xeljanz 10 mg twice daily compared to Xeljanz 5 mg twice daily or TNF blockers.

#### *Appendix D: General Information*

- Definition of failure of MTX or DMARDs
  - Child-bearing age is not considered a contraindication for use of MTX. Each drug has risks in pregnancy. An educated patient and family planning would allow use of MTX in patients who have no intention of immediate pregnancy.
  - Social use of alcohol is not considered a contraindication for use of MTX. MTX may only be contraindicated if patients choose to drink over 14 units of alcohol per week. However, excessive alcohol drinking can lead to worsening of the condition, so patients who are serious about clinical response to therapy should refrain from excessive alcohol consumption.
- Examples of positive response to therapy may include, but are not limited to:
  - Reduction in joint pain/swelling/tenderness
  - Improvement in erythrocyte sedimentation rates/C-reactive protein (ESR/CRP) levels

#### Improvements in activities of daily living

- Psoriatic Arthritis: According to the 2018 American College of Rheumatology and National Psoriasis Foundation guidelines, TNF inhibitors or oral small molecules (e.g., methotrexate, sulfasalazine, cyclosporine, leflunomide, apremilast) are preferred over other biologics (e.g., interleukin-17 inhibitors or interleukin-12/23 inhibitors) for treatment-naïve disease. TNF inhibitors are also generally recommended over oral small molecules as first-line therapy unless disease is not severe, member prefers oral agents, or TNF inhibitor therapy is contraindicated

#### *Appendix E: Mayo Score*

- Mayo Score: evaluates ulcerative colitis stage, based on four parameters: stool frequency, rectal bleeding, endoscopic evaluation and Physician's global assessment. Each parameter of the score ranges from zero (normal or inactive disease) to 3 (severe activity) with an overall score of 12.

Score	Decoding
0 – 2	Remission
3 – 5	Mild activity
6 – 10	Moderate activity
>10	Severe activity

- The following may be considered for medical justification supporting inability to use an immunomodulator for ulcerative colitis:
  - Documentation of Mayo Score 6 – 12 indicative of moderate to severe ulcerative colitis.

#### *Appendix F: The 2010 ACR Classification Criteria for RA*

Add score of categories A through D; a score of  $\geq 6$  out of 10 is needed for classification of a patient as having definite RA.



**CLINICAL POLICY**  
Tofacitinib/Tofacitinib XR

<b>A</b>	<b>Joint involvement</b>	<b>Score</b>
	1 large joint	0
	2-10 large joints	1
	1-3 small joints (with or without involvement of large joints)	2
	4-10 small joints (with or without involvement of large joints)	3
	> 10 joints (at least one small joint)	5
<b>B</b>	<b>Serology (at least one test result is needed for classification)</b>	
	Negative rheumatoid factor (RF) <i>and</i> negative anti-citrullinated protein antibody (ACPA)	0
	Low positive RF <i>or</i> low positive ACPA <i>* Low: &lt; 3 x upper limit of normal</i>	2
	High positive RF <i>or</i> high positive ACPA <i>* High: ≥ 3 x upper limit of normal</i>	3
<b>C</b>	<b>Acute phase reactants (at least one test result is needed for classification)</b>	
	Normal C-reactive protein (CRP) and normal erythrocyte sedimentation rate (ESR)	0
	Abnormal CRP or abnormal ESR	1
<b>D</b>	<b>Duration of symptoms</b>	
	< 6 weeks	0
	≥ 6 weeks	1

*Appendix G: Clinical Disease Activity Index (CDAI) Score*

The Clinical Disease Activity Index (CDAI) is a composite index for assessing disease activity in RA. CDAI is based on the simple summation of the count of swollen/tender joint count of 28 joints along with patient and physician global assessment on VAS (0–10 cm) Scale for estimating disease activity. The CDAI score ranges from 0 to 76.

CDAI Score	Disease state interpretation
≤ 2.8	Remission
> 2.8 to ≤ 10	Low disease activity
> 10 to ≤ 22	Moderate disease activity
> 22	High disease activity

*Appendix H: Routine Assessment of Patient Index Data 3 (RAPID3) Score*

The Routine Assessment of Patient Index Data 3 (RAPID3) is a pooled index of the three patient-reported ACR core data set measures: function, pain, and patient global estimate of status. Each of the individual measures is scored 0 – 10, and the maximum achievable score is 30.

RAPID3 Score	Disease state interpretation
≤ 3	Remission
3.1 to 6	Low disease activity
6.1 to 12	Moderate disease activity
> 12	High disease activity

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

#### Appendix I: Clinical Juvenile Arthritis Disease Activity Score based on 10 joints (cJADAS-10)

The cJADAS10 is a continuous disease activity score specific to JIA and consisting of the following three parameters totaling a maximum of 30 points:

- Physician's global assessment of disease activity measured on a 0-10 visual analog scale (VAS), where 0 = no activity and 10 = maximum activity;
- Parent global assessment of well-being measured on a 0-10 VAS, where 0 = very well and 10 = very poor;
- Count of joints with active disease to a maximum count of 10 active joints\*

\*ACR definition of active joint: presence of swelling (not due to currently inactive synovitis or to bony enlargement) or, if swelling is not present, limitation of motion accompanied by pain, tenderness, or both

cJADAS-10	Disease state interpretation
≤ 1	Inactive disease
1.1 to 2.5	Low disease activity
2.51 to 8.5	Moderate disease activity
> 8.5	High disease activity

## V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Tofacitinib immediate-release (Xeljanz)	pcJIA	<ul style="list-style-type: none"> <li>• 10 kg ≤ body weight &lt; 20 kg: 3.2 mg (3.2 mL oral solution) PO BID</li> <li>• 20 kg ≤ body weight &lt; 40 kg: 4 mg (4 mL oral solution) PO BID</li> <li>• Body weight ≥ 40 kg: 5 mg PO BID</li> </ul>	10 mg/day
	PsA	5 mg PO BID	
	RA		
	UC	<u>Induction:</u> 10 mg PO BID for 8 weeks, up to 16 weeks <u>Maintenance:</u> 5 mg PO BID	Induction: 20 mg/day  Maintenance: 10 mg/day
Tofacitinib extended-release (Xeljanz XR)	PsA	11 mg PO QD	11 mg/day
	RA		
	UC	<u>Induction:</u> 22 mg PO QD for 8 weeks, up to 16 weeks <u>Maintenance:</u> 11 mg PO QD	Induction: 22 mg/day  Maintenance: 11 mg/day

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

#### VI. Product Availability

Drug Name	Availability
Tofacitinib immediate-release (Xeljanz)	Tablets: 5 mg, 10 mg Oral solution: 1 mg/mL
Tofacitinib extended-release (Xeljanz XR)	Tablets: 11 mg, 22 mg

#### VII. References

1. Xeljanz/Xeljanz XR Prescribing Information. New York, NY: Pfizer Labs; December 2021. Available at: [www.xeljanz.com](http://www.xeljanz.com). Accessed January 31, 2023.
2. Singh JA, Furst DE, Bharat A, et al. 2012 Update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care Res.* 2012;64(5):625-639.
3. Ramiro S, Nikiphorou E, Sepriano A, et al. ASAS-EULAR recommendations for the management of axial spondyloarthritis: 2022 update. *Ann Rheum Dis.* 2023 Jan;82(1):19-34. doi: 10.1136/ard-2022-223296.
4. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis.* 2015;0:1-12. doi:10.1136/annrheumdis-2015-208337
5. Rubin DT, Ananthakrishnan AN, Siegel CA, Sauer BG, Long MD. ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol.* 2019 March;114(3):384-413. doi: 10.14309/ajg.000000000000152.
6. Singh JA, Saag KG, Bridges SL, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Rheumatology.* 2016. 68(1):1-26.
7. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for non-systemic polyarthritis, sacroiliitis, and enthesitis. *Arthritis Care & Res.* 2019; 71(6):717-734. doi: 10.1002/acr.23870.
8. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol.* 2019;114:384-413.
9. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the treatment of psoriatic arthritis. *American College of Rheumatology.* 2019; 71(1):5-32. doi: 10.1002/art.40726.
10. Smolen JS, Landewe RB, Dergstra SA, et al. 2022 update of the EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs. *Arthritis Rheumatology.* 2023 January; 32:3-18. DOI:10.1136/ard-2022-223356

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Per HFS PDL, new policy created, adapted from CP.PHAR.267 Tofacitinib (Xeljanz Xeljanz XR) policy.	12.13.21	
2Q 2023 annual review: no significant changes; updated off-label dosing in Appendix B; Template changes applied to other	4.15.23	

**CLINICAL POLICY**  
**Tofacitinib/Tofacitinib XR**

Reviews, Revisions, and Approvals	Date	P&T Approval Date
diagnoses/indications and continued therapy section; updated section I initial criteria; references reviewed and updated.		

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

## CLINICAL POLICY

### Tofacitinib/Tofacitinib XR

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.